

THE MEDICAL NEWS.

A WEEKLY JOURNAL OF MEDICAL SCIENCE.

VOL. LVI.

SATURDAY, JUNE 28, 1890.

No. 26.

ORIGINAL LECTURES.

ENUCLEATION OF GOITRE.

*A Clinical Lecture
delivered at the Jefferson Medical College Hospital,
March 22, 1890.*

BY W. W. KEEN, M.D.,
PROFESSOR OF SURGERY.

GENTLEMEN: I shall show you to-day a case of goitre in a woman with the following history: She is thirty years of age, an American by birth. Six years ago she noticed that breathing after exertion became difficult, and three months later she discovered a slight swelling in the neck. This swelling has increased in size only gradually, except for two or three months after each of her two labors, when the increase was quite rapid. This statement is in consonance with the fact that in many women there is increased fullness of the thyroid gland after maternity. The tumor is not painful. The urine is straw-colored and contains a doubtful trace of sugar, some pavement epithelium, but no albumin.

The following measurements have been taken: Circumference of the neck $16\frac{1}{4}$ inches. The horizontal arc of the tumor, from side to side, is $7\frac{3}{4}$ inches, subtending a chord of $4\frac{1}{4}$ inches. The horizontal arc from the middle line to the right border is $4\frac{1}{2}$ inches; from the middle line to the left border $3\frac{1}{4}$ inches. The vertical arc is 6 inches; its chord $3\frac{1}{4}$ inches. The tumor feels distinctly cystic and resilient, but it is not markedly lobulated. The larynx has been examined by Dr. Jones and the vocal cords found to be normal, and there is no paralysis. I should add that it is through the courtesy of Dr. Hearn that I have the opportunity of showing you this patient.

There are three kinds of goitre: First, we have exophthalmic goitre, which heretofore has been a non-surgical affection, and is accompanied with enlargement of the heart, rapid pulse and protuberance of the eyeballs. This form has been successfully operated on recently, but the number of cases is too small to generalize from. Second, we have parenchymatous goitre, in which the gland tissue proper of the organ is involved. This makes a hard dense fibrous structure. Third, we have the form which I think is present in this case—a cystic goitre, which is really a fibro-adenoma with one or more cysts. In addition to these goitres, properly so called, we have the various cancerous enlargements of the gland, which are necessarily more serious than those which I have mentioned.

The diagnosis of goitre is usually easily made, but the variety is not always so readily determined before operation.

The medical treatment of goitre is unsatisfactory. Sorbefacient and irritant ointments and internal remedies have given practically but little result. In the surgical

treatment of this affection several methods have been employed. The first method that I shall allude to is that of injection, either of the ordinary tincture of iodine, or as proposed by Sir Morell Mackenzie, of the tincture of the chloride of iron, the object being to convert the enlargement into an abscess. I do not believe in either of these methods, especially since the splendid results of operations which have been reported by Socin and others. In the light of these reports, I think that it is useless to talk of the injection of any fluid into these tumors. It is a procedure attended with danger of septic infection, and is painful, tedious, and uncertain in its results.

Of operative procedures proper, there are three. The first is evacuation. In some cases, especially of large immovable masses, this will do a great deal of good. It consists in opening the various loculi, whether cystic or solid, and peeling out their contents. The second method, and the one which I shall try in this case, is enucleation. It consists in cutting down to the cyst itself and enucleating it as we would any other cystic growth. This method has been advocated especially by Socin, of Switzerland, and his cases have been reported by Koser, his assistant. The third method is extirpation, which is the most dangerous of all, both as concerns its immediate and subsequent results. In 1871 Dr. Green, of Portland, Maine, reported six cases of this kind, and it surprised the surgical world that he was able to report so many. Since that time, and more especially in the last few years, a revolution has taken place in the surgery of the thyroid gland. Billroth has reported a large number of cases in which he has had remarkable results. It has remained for Kocher, of Switzerland, to give the most extraordinary figures. In a paper published in the *Correspondenzblatt für Schweizer Aerzte*, January, 1889, he reports 250 cases of extirpation. His results are astonishing. Of the 250 cases 25 were cases of cancer. Of this number 4 died, a mortality of 17 per cent. Of the remaining 225 non-cancerous cases, 223 recovered and only 2 died, a mortality of only 0.8 per cent. Socin in 59 cases of evacuation and enucleation had only one death. This occurred a short time after operation and was due to hæmorrhage. You can imagine how great the hæmorrhage was, when I tell you that during the operation 170 ligatures were employed. Years ago I saw a case of goitre operated on in this city in which the hæmorrhage was enormous, and the patient died three days afterward.

I have here some drawings from Kocher's paper published in Langenbeck's *Archiv* for 1880. The incision which he recommends in extirpation is an angular one, vertical in the median line and then extending outward over the larger portion of the growth. If necessary, this can be converted into a Y-shaped incision by an oblique cut on the opposite side. In operating it is the veins which give by far the greatest trouble. In this first in-

cision the external jugular will probably not be interfered with, but the anterior jugular, another vein called by Kocher the oblique jugular, and the anterior communicating branch will be cut. These are to be tied with double ligatures and the vein divided between them. We next come to the deeper vessels. We have a superior communicating vein going from the superior thyroid vein of one side to that of the other. There is also a deep anterior communication connecting the deep thyroid veins of each side. There is not only a superior thyroid, but also an accessory superior thyroid. In the same way there is an inferior thyroid and an accessory inferior thyroid. Then we have, what is very common, a still lower thyroid vein, the *vena thyroidea ima*, with an accessory vein to it. These veins are to be divided methodically. The order in which the three superficial veins are tied makes little difference. The fourth vessel to be tied is the superior thyroid; fifth, the accessory superior thyroid; sixth, the superior communicating; seventh, the *thyroidea ima*; eighth, its accessory; ninth, the inferior thyroid; tenth, its accessory; and eleventh, the inferior communicating.

But there are also arteries as well as veins which require attention. It is principally at the inferior thyroid that trouble is apt to be met with. This is because of its relation with the recurrent laryngeal nerve. The inferior thyroid, as you know, arises from the thyroid axis, goes behind the carotid and the sympathetic nerve, having in front of it the superior ganglion of the sympathetic, from which goes the cardiac nerve. The artery then goes behind the recurrent laryngeal nerve and passing under the tumor toward the middle line, finally plunges into the gland. In tying this artery there is danger of wounding the sympathetic or the recurrent laryngeal nerves. If the latter is destroyed it produces paralysis of the vocal cord of that side, but does not necessitate tracheotomy. If both are injured it becomes necessary to do tracheotomy, as the glottis closes.

The only way to avoid this injury to the nerve is to *isolate absolutely* the vessel before it is tied. In goitre this artery never seldom becomes as large as the carotid.

Another difficulty is that the recurrent laryngeal nerve is very tightly adherent to the gland on its posterior surface in the groove between the œsophagus and trachea. If the gland is drawn over to one side, the nerve may be pulled into a loop and injured, causing paralysis of the vocal cord. It is important to keep close to the tumor at this point.

I have here some statistics taken from Keser's paper, published in 1887. The number of cases operated on was 59. Of these, 5 underwent total extirpation, 17 partial extirpation and 37 enucleation. It was found that in 9 of the partial extirpations, enucleation could have been performed. The ages of the 59 patients varied between twelve and seventy years. There was only one death. In 11 partial extirpations, the histories of which were followed, there was return in 5. In 21 enucleations there was return in only 3.

The dangers of extirpation are immediate (as from hæmorrhage) and remote. The remote dangers are important. These dangers have been explained by the experiments of Horsley. In cases of atrophy or destruction of the thyroid gland its function is, of course, abolished. The function of this gland seems to be a

two-fold one. It presides in some obscure way over the nutrition of the mucous tissues of the body, and in the second place it largely controls the genesis of the blood. The complete extirpation of the gland is liable to be followed by great anæmia and a condition which is called myxœdema or cachexia strumipriva. In this curious condition the body becomes bloated, the eyelids hang in folds, the face is like that of a person dropsical from Bright's disease, but the exudation is not soft but hard, and results from the development of mucous tissue. After a time, the hair drops out and mental changes occur, finally resulting in crétinism. This is often seen in Switzerland in connection with goitre. In these cases the speech is slow and faulty, mental operations become sluggish, albumin appears in the urine, and finally death closes the scene. Another curious occasional result is that peculiar muscular tremor known as tetany. Myxœdema and tetany are not apt to occur in those cases in which a portion of the gland is allowed to remain. This fact has led to enucleation of the goitre in preference to extirpation of the whole gland, and lately to the recommendation by some that a part of the gland should be transplanted. This has been suggested by Schiff and Kocher, and has recently received the approval of Horsley. The operation consists in taking a part of the normal thyroid gland of some animal, as a sheep, and transplanting it into some part of the body before operation, as, for instance, into the abdominal cavity. The results of these operations have not yet been published, but, as I have stated, they have received the approval of so able a surgeon and experimenter as Horsley, which is a great point in their favor.

In this case I propose, if possible, to do enucleation and not extirpation. In enucleation we avoid two of the dangers of extirpation, the immediate danger of injury to the recurrent laryngeal nerve and the remote danger of myxœdema or tetany. Here a large portion of the left lobe appears to be in good condition, and I shall try to preserve it.

In some cases it is necessary to do tracheotomy, but always avoid it if possible, for the moment you open the trachea you have a septic wound, as all the air entering and passing from the lungs passes over the wound.

You observe the peculiar breathing which the patient shows, and also the fact that she does not take ether well. I first make my incision through the skin and ligate all the veins that appear. The layer of gland tissue which overlies the cyst in these cases may be very thin or it may be as thick as one-half or three-fourths of an inch. I shall cut through this tissue until I reach the cyst. I now come to what I think is the cyst wall. I ligate the veins that appear on its surface, tying every vessel that would be apt to bleed. I now find, however, that it is not the cyst wall, but glandular tissue, almost one-fourth of an inch of which I must cut through before I reach the cyst. On opening this, a quantity of dark-colored fluid is discharged. It is easily seen by those who are near that the cyst-wall can be readily distinguished from the gland tissue. The cyst, as you see, is separated and removed without difficulty, with but little bleeding. There is quite free hæmorrhage from one rather large vessel at the bottom of the cavity, but this is now controlled after some little trouble. I shall not close the wound until satisfied that every bleeding vessel

is tied. The number of ligatures used by Socin varied between twenty-four and sixty. In this case I have used thirty-seven. I have not at all interfered with the left lobe of the gland, which, however, is only slightly enlarged. As there is no further bleeding I apply deep catgut sutures to bring together first the cyst wall, then the muscles, and lastly the skin, by superficial sutures of silk, inserting a rubber drainage-tube and also a bundle of horsehair. The wound is then dressed in the ordinary manner.

NOTE.—The patient had had a troublesome cough for some days before the operation. This was very markedly increased immediately after the operation. She was also uncontrollably restless, and so displaced the dressing that the wound was exposed and became infected within twenty-four hours; moderate suppuration and considerable fever followed, the temperature rising to 104.2°. For six days she was seriously ill. The sutures in the vertical incision were removed, the wound washed out with bichloride solution and ample drainage secured. Alcohol was freely given. She then began to improve; in ten days the temperature became normal, and she left the hospital, after three weeks, entirely recovered.

ORIGINAL ARTICLES.

SUPRA-VAGINAL HYSTERECTOMY.


Hysteromyomectomy with Suspension of the Stump in the Lower Angle of the Abdominal Incision.

BY HOWARD A. KELLY, M.D.,

GYNCOLOGIST TO THE JOHNS HOPKINS HOSPITAL, BALTIMORE.

THE method here proposed¹ is limited to cases conceded to be most favorable for operation—those in which there naturally exists a pedicle, or in which it is possible to form a pedicle below the tumor masses. I do not wish to consider atypical cases, in which total ablation of the uterus (panhysterectomy) is called for, or those in which the uterus

FIG. 1.




TTT, tumor. P, pedicle. V, vagina in section.

from fundus to cervix is a mass of fibroid tumors. I also purposely avoid the important question as to the best method of forming a pedicle when the

broad ligament is choked with fibroid tumors. Fig. 1 is a generic representation of the class under discussion.

I have adopted in these cases an original method combining the advantages and eliminating many of the dangers of Hegar's and Schröder's methods—the ordinary extra- and intra-peritoneal methods.

FIG. 2.



The operation consists of seven steps, as follows: First. A long incision in the linea alba (Fig. 2) for the delivery of the myomatous uterus.

Second. The elevation of the tumor until the pedicle is brought into view for treatment by tying the broad ligament structures, or for enucleation of tumors from the broad ligament until a pedicle is formed, when the rubber ligature is applied and tied tightly, controlling the circulation. (Fig. 3.)

Third. The tumor is cut away from one to two inches above the rubber ligature (Fig. 4), by first

FIG. 3.



Tumor delivered. rr, rubber ligature in place and ready to be drawn tight. Dotted line shows the incision through the peritoneum.

splitting the peritoneum and then cupping out the upper face of the stump, cutting with each stroke down toward the vaginal canal.


The cervical canal must next be carefully dissected out and its site well cauterized.

Fourth. This consists in the closure of the raw face of the stump by uniting the opposite sides by means of a continuous buried suture of catgut as seen in Figs. 5 and 6. Fig. 5 represents the appearance

¹ See American Journal of Obstetrics, April, 1889.

seen upon looking down on the stump from above, Fig. 6 being a vertical section through the cervix.

FIG. 4.




Tumor separated from pedicle.

FIG. 5.



Method of closing upper raw surface of stump by means of a continuous buried suture.

FIG. 6.




Stump united up to last row of interrupted surface-sutures, *SS*. *bs*, last rows of buried sutures. *V*, vagina. *cv*, vaginal cervix.

The last row of sutures, which brings the peritoneal surfaces into apposition, is of interrupted silk sutures, with the long ends left uncut, for a purpose to be described later. All of these sutures, buried and superficial, must be applied with the view of controlling the circulation as well as securing approximation. They must, therefore, be drawn tight, and must encircle any vessels in view.

Fifth. After the surface of the stump has thus been closed and there remains nothing of the wound but the linear union of the peritoneal surfaces, the rubber ligature is cut away, and the lips of the wound are carefully observed. If there is any persistent oozing the nearest uterine artery must be ligated. This is accomplished by grasping the long ligatures and pulling the stump to the right or left, exposing the site of the left or right artery. A stout needle armed with a catgut ligature is then swept boldly through the side of the stump, well below the sutured area (Fig. 7) and tied, thus cut-

FIG. 7.



Ligation of left uterine artery. Pedicle is pulled toward the right by means of the long interrupted silk sutures *SS*, while the needle carrying ligature is passed under the uterine artery.


ting off all communication between the artery and the stump.

Both arteries may be treated in this way without any danger of destroying the vitality of the stump. If, however, there should be no flow of blood from the closed lips of the stump, the fifth step may be omitted.

Sixth. The abdominal incision is closed down to

the stump, putting in a drainage-tube, if needed, well above the stump. Following this the parietal peritoneum of the abdominal incision is united to the peritoneal coat of the stump, below the lips of the stump, by means of a continuous catgut or silk ligature (Fig. 8); and in this way the stump is separated from the peritoneal cavity.


FIG. 8.



Showing union of the peritoneal surface of the stump to the parietal peritoneum.

Seventh. The wound is dressed with some dry antiseptic powder, or simply packed under the edges of the skin, around the suspended stump, with antiseptic gauze. Finally, a large square of gauze, six or eight folds in thickness, with a small slit in it, is prepared, and the long ligatures which unite the peritoneal lips of the stump are pulled through the slit. These are lifted well up, and grasped by a pair of long Keith's forceps laid horizontally on the body (Fig. 9).

FIG. 9.



Dressing applied. Interrupted sutures of the surface of stump are brought through a hole in the gauze and grasped by forceps.


This dressing serves effectually to keep the stump from pulling back into the abdomen, and the operator has at all times full control of it at a moment's notice in case of accident. This gauze can be

changed as often as soiled. Once every two or three days is usually sufficient. The silk sutures uniting the peritoneal lips of the wound finally either come away or are cut loose and pulled out in about ten days.

The small pit which is thus left in the lower angle of the abdominal wound over the stump rapidly fills by granulation.

This method has distinct advantages over internal or external methods commonly in use. It is better than dropping the stump back among the intestines (Fig. 10). By the new method hæmorrhage is not

FIG. 10.



Intra-peritoneal treatment. Stump *S* dropped back into peritoneal cavity and abdominal walls closed above.

dangerous, being at all times under control. The danger of sepsis is also removed, a danger to which large numbers of cases have succumbed after the intra-peritoneal treatment of the stump.

It is better than the common external method (Fig. 11), because, in the first place, it is there

FIG. 11.



Extra-peritoneal treatment of stump *S*, which sloughs off at the rubber ligature *r*. The union of the peritoneum of the stump to that of the abdominal walls is shown by the dotted circles.

necessary to elevate even a short pedicle far enough to attach the parietal peritoneum *below the rubber ligature*. By my method the attachment is higher, and the method is, therefore, better for short pedicles, doing away with a traction which is often excessive.

Again, when the rubber ligature is left on, it is impossible to limit the depth of the slough which takes place as the distal end drops off, and it will readily be granted on general principles that a method which constricts any part of the body, and waits for it to drop off by sloughing is a coarse and unscientific means of performing an amputation.

THE PROPRIETY OF THE REMOVAL OF THE APPENDIX VERMIFORMIS DURING THE INTERVALS OF RECURRENT ATTACKS OF APPENDICITIS.¹

BY FREDERIC S. DENNIS, M.D.,

VISITING SURGEON TO BELLEVUE AND TO ST. VINCENT'S HOSPITALS,
NEW YORK.

THE question propounded in the title of this paper is a new one in surgery. Its solution cannot be made at once. The entire subject must first be considered in the light of aseptic surgery, and, having discussed the question from that point of view, the operation must be examined into as to its propriety in certain cases. A definite answer must be postponed until new data and recent clinical experience have afforded us more reliable information from which to judge the expediency of such an important surgical procedure. The propriety of excision of the appendix can be discussed, however, from a theoretical point of view, and reference can also be made to the few reported cases, and from such data some knowledge can be obtained which will at least throw light upon this hitherto unexplored field of operative surgery. The entire subject of iliac inflammations is fraught with great interest from many points of view. However attractive these inflammations may appear, the discussion in this paper is restricted to one single question in regard to the whole subject, and that is, Is the surgeon justified in excising the appendix after recovery from one or many attacks? The question of removal of the appendix during an interval of quiescence cannot be answered dogmatically. I shall endeavor to demonstrate that the operation as routine treatment in all recurrent cases is unsurgical, and that only in rare cases is it indicated.

With the view to simplifying the subject it seems pertinent to classify these inflammatory attacks; because all inflammations in this region are not identical, either in etiology, in the anatomical part involved, or in the termination.

The opinion has recently been advanced that all of these conditions are dependent upon a primary pathological change in the appendix, and that all the subsequent changes are secondary to an affection of the appendix. For my own part I cannot endorse this view. I have observed many cases

that certainly ran a course very similar to typical appendicitis, and yet the cæcum or the pericæcal region were the parts involved, and the appendix was normal. I shall have occasion to refer again to these cases. If the view is accepted that all these changes are secondary to a primary appendicitis, then certainly it must be admitted that this one region forms an exception to the laws governing inflammations in other parts. This belief would imply that there is a part or section of the alimentary canal that enjoys an immunity from primary inflammatory changes, and that there is a limited area of connective tissue in the right iliac region which is free from initial inflammations, and that there is a restricted zone of peritoneum exempt from a peritonitis except as secondary to an appendicitis. Surely this view is not tenable, and clinical experience clearly points to the conclusion that there is no exception here to the laws governing inflammations in general; but that each part, and every structure, and all varieties of tissue may become the seat of an inflammation independent of an appendicitis. To refer now to a classification of these inflammations. They may, for convenience, be divided into—

1. Typhlitis.
1. Perityphlitis.
3. Appendicitis.
4. Peri-appendicitis.

These different inflammations may merge into one another if no surgical treatment intervenes; but the separate conditions may exist independently, and each may run its course without necessarily involving the other. By a typhlitis is meant an acute or subacute inflammation of one of the several coats of the cæcum. This form may possibly extend to the adjacent tissues. The lesion consists of a catarrhal or ulcerative inflammation, and begins in the mucous membrane. This variety of inflammation is dependent upon faecal impaction, a septic diarrhoea, a corrosive poison, or an ulcerative process.

There are some surgeons who deny the existence of a localized inflammation of the cæcum. I am confident that I have seen several cases of this kind. The proof of this statement becomes apparent by an examination of two specimens which I have, showing well-marked ulceration of the mucous membrane of the cæcum, and these changes are entirely independent of any trouble with the appendix. Dr. Janeway informs me that he has recently seen three cases of localized cæcal inflammation. These cases certainly establish the clinical fact which is denied by some surgeons, that a local inflammation in the right iliac region can exist independent of appendicitis.

The case reported by Mr. Pepper, of London, is another proof of local cæcal inflammation. The vaginal injection of a 1:2000 bichloride of mer-

¹ Read before the American Surgical Association, May 14, 1890.

cury solution caused extensive ulceration of the cæcum. The patient died, and the autopsy revealed the fact that the ulceration was limited to the cæcum.

Toxic enteritis may chiefly, if not entirely, involve the cæcum, and after the administration of antimony as well as of arsenic this phenomenon has followed. I have cured a case of extensive ulceration, of eleven years' standing, of the cæcum and ascending colon, by copious injection high into the colon, using as strong a solution of bichloride as the circumstances permitted. After eleven years of septic diarrhoea and hæmorrhage this patient was entirely cured of his cæcal and colon ulceration.

The second variety, viz., perityphlitis, is that form of inflammation which involves the areolar tissue in the vicinity of the cæcum. This leads to suppuration and may exist independent of any change in the appendix. These cases are observed following traumatism, as in sprains of the psoas muscle, or from some pathological condition in the ilium, sacrum, or vertebrae. I have seen these cases occur after violent athletic exercise; one case occurred after swimming, a second after jumping, and a third after lifting heavy weights. These abscesses go on to free suppuration, are situated in a kind of tissue prone to break down after an injury, are not subject to perforation, but generally point toward the periphery of the body.

The third variety, appendicitis, means an inflammation of the vermiform appendix, and is identical with a typical typhlitis, only that it is more serious in its consequences, owing to the anatomical formation of the part; but the causes which give origin to the change are, in many respects, similar.

The fourth variety, viz., peri-appendicitis, is not so clearly defined as is perityphlitis, because this variety is undoubtedly secondary to an appendicitis. The small amount of areolar tissue about the tip of the appendix is not exposed to the same injuries as the more abundant quantity about the cæcum.

The question of removing the appendix during the quiescent period would therefore involve the last two conditions, viz., an appendicitis or a peri-appendicitis.

It is, then, with an appendicitis or a peri-appendicitis that this question of removal of the organ during an interval of attack concerns us at the present time.

In discussing the operation the first question that naturally arises is, What proportion of cases of appendicitis are recurrent? It is with this percentage of cases that the question of removal of the appendix arises. There are no definite statistics upon this most important point. Fitz, in his scholarly article, has found in the examination of 257 cases of appendicitis that 28 of the cases were

recurrent. This would place the number at about 11 per cent. In other words, 89 per cent. of the cases of appendicitis are solitary and 11 per cent. are multiple. The proportion is about the same in regard to typhlitis and perityphlitis as regards solitary and multiple attacks.

A second question now arises, viz., In the 11 per cent. of cases in which a relapse occurs, what is the termination? If all the 11 per cent. of cases of relapsing appendicitis terminated fatally then the question of excision of the appendix is easy to answer. It is obvious that if all these patients died during one of the recurrent attacks excision of the appendix should be attempted after the first or second attack; because the removal of the appendix would be the only salvation open to these unfortunate individuals. Happily death is not the only termination for this 11 per cent. of cases.

They may terminate in:

1. Resolution.
2. Formation of pus and evacuation of abscess.
3. Perforation and general peritonitis.

It is not known how many of these cases of relapsing appendicitis terminate in resolution. Unfortunately there are no statistics upon this point.

Nearly all of the authorities, however, are in accord with the general statement that a majority of the cases undergo resolution, and that only exceptionally perforation occurs.

The second way in which an appendicitis may terminate is by the formation of pus and the evacuation of the abscess, either spontaneously or by surgical interference. In regard to the chances of recovery offered by an early operation, it has been shown that there are fifty per cent. more recoveries following operations before the eighth day of the disease than after the eighth day, and that the mortality is only eight and one-half per cent. in those cases operated upon before the eighth day. If now the operation of incision is performed during the second or third day of the disease in these relapsing cases, the rate of mortality is lessened to such a degree (as shown by a study of Weir's, Bull's, and McBurney's cases) that the mortality is comparatively insignificant.

If now out of the eleven per cent. of cases of relapsing appendicitis there are deducted first those cases which terminate by resolution, and second, those cases which recover by early incision, it is obvious that only a small part of the eleven per cent. of cases are exposed to the remaining source of danger, viz., perforation and general peritonitis.

The whole question of removal of the appendix, therefore, practically has reference only to those cases in which a general peritonitis, the result of a perforation, is likely to arise. It is under these circumstances that death follows, and the excision of the appendix is the only measure that will remove

this cause of almost certain death. The line of argument thus pursued brings us face to face with the clinical fact that removal of the appendix is to be considered only as a means to protect a patient in the future from death by a peritonitis the result of a perforation. If it can now be shown that in the extreme emergency of a general peritonitis a laparotomy can save a patient, then the removal of the appendix during a quiescent period is deprived in a great measure of the only argument that can be claimed for its performance. Simple reference to the cases of Krönlein, Morton, Weir, Sands, McBurney, and others demonstrates that perforation does not necessarily preclude all possibility of recovery, which is the great argument in defence of removal of the appendix during the period of quiescence. The same line of argument may be carried even further, and it can be shown that perforation may occur and no laparotomy be performed, with still a possible chance for recovery. A proof of this statement may be found by a reference to the following cases.

Dr. Markoe reports the case of a child who had symptoms of general peritonitis upon the second day of an attack. A month later this child died from some other disease and the autopsy showed that perforation of the appendix had occurred, and that adhesions of the intestine had taken place. Dr. Biggs made an autopsy and found the evidences of a previous perforation of the appendix from sloughing. The appendix was found encysted in an old circumscribed peritoneal abscess.

A case is mentioned by Shaw where an abscess was formed in the scrotum from a perforated hernial appendix, and Thurman records a similar case.

Dr. Cabot reports a case where an inch of a previously perforated appendix was found in an abscess in the groin some eighteen months after an attack of appendicitis.

Ballou and Pooley each report a case where the appendix had sloughed and the patients recovered. Many other cases might be cited to show that even if perforation of the appendix has occurred it is not necessarily a fatal lesion.

Dr. Janeway made an autopsy on a patient who died some time after an attack of appendicitis. The appendix had been the seat of ulceration and perforation; but it had formed adhesions by inflammation to the surrounding tissues and no gas had escaped into the general peritoneal cavity, and as far as the perforation was concerned, the patient lived. It is an illustration of the clinical fact that a circumscribed peritonitis may shut off the appendix from the general peritoneal cavity and that even perforation may occur and not be a fatal lesion.

The best of all proofs that perforation of the appendix may occur and the patient survive such an

attack, is found in a case in which an autopsy was made ten years after an attack of appendicitis. It was ascertained that the appendix had been perforated during the attack and that it had been accompanied with abscess and suppuration. Dr. Biggs, who made the autopsy, found that the tip of the appendix had sloughed from its base, to which it was still held by an impervious cord. The entire mass, including the base and apex, with the filiform connection between the two separated parts, was embedded in a mass of inflammatory new-formation tissue, which was adherent to the kidney and to the liver. This patient for ten years had no trouble with his appendix, and no one ever suspected the nature of his previous attack of illness.

If now I have made the thread of my argument clear, it is obvious that excision of the appendix has to do with only about 11 per cent. of the cases of appendicitis, for 89 per cent. of the cases are not relapsing, and therefore do not belong to the class of cases in which removal is indicated. Of the 11 per cent. a majority recover; because perforation is the exception rather than the rule. Some undergo suppuration and heal. Of the few cases exposed to the danger of perforation and general peritonitis some recover, as is shown by a reference to the cases just mentioned.

It is now pertinent to consider in detail the real arguments against removing the appendix during a quiescent period in the few cases where the attacks have been recurrent. The arguments which I would respectfully submit against this operation of excision during a quiescent period in relapsing cases are as follows:

1. The danger to human life.
2. The difficulties of a positive diagnosis.
3. The development of ventral hernia.
4. The lack of conclusive evidence that excision of the appendix is attended with permanent relief.
5. The result of relapsing attacks may afford an immunity from danger in the future.

1. *The danger to human life* is an argument which confronts the conscientious surgeon. Opening the abdomen for the purpose of finding the appendix vermiformis is certainly attended with risk to life. That risk increases in accordance with the amount of pathological change that has taken place. I must dissent from the view held by Morton, who speaks of this procedure as a "comparatively trivial operation at a time and under conditions when prompt and permanent relief and recovery can almost invariably be secured." The same opinion is also expressed by Senn, who writes: "Excision of the appendix in cases of simple uncomplicated appendicitis is one of the easiest and safest of all intra-abdominal operations." Such statements are misleading. In May, 1887, I excised the appendix, on

account of a stab-wound, with no difficulty as regards finding it; but to remove it and to invert the edges and to introduce sutures and to close hermetically the opening into the cæcum is a delicate operation which requires nicety of *technique*. It is by no means to be considered a trivial procedure, and upon the result of the operative work of the surgeon depends the life of the patient. The giving way of a stitch or the sloughing of a small shred exposes the patient to imminent peril. One bubble of intestinal gas is sufficient to infect the whole peritoneum, and such a catastrophe will in all probability lead to a fatal termination. The possibilities of hæmorrhage, of purulent œdema, of septic peritonitis, of suppression of urine, and of surgical shock must not be overlooked. These are the dangers to which a patient is subjected in the ordinary cases; but when, from recurrent attacks, adhesions have formed, the connective tissue has undergone pathological changes, the cæcum has become dilated, and the anatomical relations are disturbed, excision of the appendix becomes a serious operation. Surgeons have attempted to find an analogy between removal of the appendix and of the ovary. This comparison causes a tendency to attach too little importance to the excision of the appendix. It must be borne in mind that the vermiform is an integral part of the alimentary canal; it often is adherent to the large and important bloodvessels, it may be buried under the cæcum, it may be firmly imbedded in inflammatory exudation, the tearing of which may result in a rent in the intestine or ureter, and the sewing up of which requires time and skill. Not so with the ovary, for here the oozing is easily controlled, and if a pus cavity is found the cavity can be easily washed out and rendered aseptic. For these reasons the excision of the appendix cannot be looked upon in any light other than that of an operation attended with considerable risk to human life.

The operation of removal of the appendix during an interval between the attacks has been attended with a death-rate the precise percentage of which cannot be at present ascertained, because a report of all the cases has not yet appeared in print. It is safe to state, however, that the death-rate is higher than that of early incision on the second or third day of an attack. This one argument alone deprives the operation of its greatest advantage. Thus far, in dealing with this subject the ability to find and to remove the appendix has been assumed; but a most serious aspect of this question is the failure to find the appendix after the abdomen has been opened, or, owing to extensive adhesions, an inability to remove the organ. Recently I was present at an operation for the purpose of excision of the appendix. The laparotomy was

performed upon the patient during the quiescent stage, and although the surgeon was one of the most skilful operators and one of the best anatomists, he was unable to find the appendix after a long and tedious search. The abandonment of this operation was wise and prudent, and I mention this clinical fact to impress the point that the operation is, of necessity, attended with great danger where a prolonged search must be made with the peritoneal cavity opened and the intestine exposed. When an operation becomes necessary it should be performed early during an attack, and at the same time removing, if necessary, the appendix. It is the proper line of treatment. It is quite another thing after an attack has passed and in the quiescent stage to attempt an excision. In the former case it is indicated to save life, in the latter case it is fraught with great danger to avoid a possible future attack, from which the patient may even easily recover.

2. *The difficulties of a positive diagnosis.*—No surgeon who has seen many cases of abdominal surgery can overlook the fact that the diagnosis is often fraught with the greatest difficulties and with much uncertainty. The more cases examined the more real becomes this fact. It is often impossible to arrive at a positive and clear diagnosis as to the real condition that has given rise to the attacks. The most skilled diagnosticians and the most experienced surgeons have made errors in diagnosis as regards lesions in this region. It is only necessary to review the list of diseases that already have been mistaken for appendicitis in order to estimate the weight of this argument against excision of the appendix during a quiescent period.

Among the conditions that have been mistaken for appendicitis may be mentioned the following: General or circumscribed peritonitis, pelvic peritonitis, renal, biliary, and intestinal colic, ovarian and lumbo-abdominal neuralgia, intestinal obstruction, floating kidney, pyelitis, cæcitis, internal strangulation, psoas abscess, pelvic cellulitis, rupture of the serratus magnus muscle, suppurative adenitis, typhoid, tubercular and stercoral ulcers, caries of ilium and of vertebræ, morbus coxarius, suppuration in the retro-peritoneal and mesenteric glands, traumatic rupture of intestine and right ureter, rupture of the bladder and of the gall-bladder, rupture of an aneurism in the broad ligament, sprain of the iliacus and psoas muscles, salpingitis of the right tube, abscess of liver, tubal pregnancy, typhlitis and perityphlitis.

This list does not embrace nearly all the conditions which eminent men in the profession have frankly acknowledged to have mistaken for appendicitis, or *vice versa*. The list, however, is of sufficient size to impress the important point that errors in diagnosis may occur, and that a laparotomy for

excision of the appendix may be performed and the real pathological condition which the operation is intended to relieve be other than a diseased appendix. Finally, it is significant that in some of the cases in which the appendix has been removed during the quiescent period no change sufficient to cause trouble could be found in the appendix. Within a month I have examined a specimen showing a perforation of the cæcum where appendicitis was diagnosed. If this patient had survived the attack and the appendix had been removed the operation would have been of no avail, for the lesion was situated in the cæcum.

3. *The development of ventral hernia.*—This interesting observation has been made by Dr. Bull, whose brilliant work in abdominal surgery is of world-wide reputation and entitles his opinion to the highest respect.

This condition is not likely to follow the early incision which is made during an attack to allow pus to escape, because the incision under these circumstances is very small, whereas, in a laparotomy for excision of the appendix during the interval of attacks, the length of the incision is from four to five inches. Several cases of ventral hernia following operations for removal of the appendix have been reported. The special situation of this incision and the peculiar character of the parts divided, and the tendency of the wound to gape, render the development of hernia a most serious complication. There is no doubt that this condition can give rise to strangulation, and even if strangulation does not follow, the presence of the ventral hernia is a source of great discomfort and annoyance to a patient.

4. *The lack of conclusive evidence that the excision of the appendix is attended with permanent relief.*—Time is the crucial test to settle this important point. The cases that have been published bear witness to the fact that the operation has been performed, but the period of time is too short to base any conclusions as to the ultimate result. If the excision has been performed in a case where the appendix alone was involved, the result may be satisfactory as regards the appendix; the pathological process, however, which established the lesion in the appendix may subsequently develop in the cæcum and changes similar to the original attack may then ensue. If the appendix has been removed when the lesion is in the cæcum the relief does not follow.

To what extent the operation will afford immunity in the future is still a question *sub judice*. There is positive evidence that recurrent attacks may terminate in a final one with recovery, and the patient enjoy years of immunity. He may never have any return of the trouble. I know of one patient residing in a southern State who has had fourteen con-

secutive attacks of appendicitis; but during the past five years has never had the slightest indication of any disturbance. These fourteen attacks occurred during a period of a few years, and the patient was able to trace the cause of the attacks to the use of cathartics and to a peculiar diet. Since he has abandoned cathartics and has lived upon a meat and gluten diet he has never had the slightest indication of trouble. His appendix may be so situated as to never develop another attack. An attempt to excise his appendix might be attended with a fatal issue.

5. *A condition following relapsing attacks may afford immunity from danger in the future.*—I feel certain that some surgeons will dissent from this view; but I shall endeavor to prove the correctness of the statement from pathological material and from clinical data. One great mistake that surgeons commit in discussing this subject is to consider all cases of appendicitis doomed to immediate death. Investigation and observations are made at the time of a contemplated incision to avert perforation, consequently the subject is viewed from very narrow limits. I believe that often a single attack from which recovery follows places the appendix in such a position that no further harm can result to the patient. For example, I have four microscopical specimens showing that the appendix in a cross-section has become completely obliterated as a result of appendicitis, and that instead of the normal duct with its narrow lumen filled with glairy mucus an impervious cord exists. This change is the result of a previous inflammatory process, as is seen by an examination of the tissue. I know of other cases where the same condition is present. If this is true in regard to a single attack, why may not the last of recurrent attacks place the appendix in this same condition? An appendix, then, that is an impervious cord becomes so by inflammatory processes, and I fail to find a report of an autopsy where the patient died from an attack of appendicitis and in which the appendix was an impervious cord.

These cases of obliterated appendices are found incidentally, and upon microscopical examination the fact is revealed that the cords are the result of a previous attack of appendicitis. But the impervious cord is not the only condition that may afford immunity to a patient. For example, I have specimens showing some interesting conditions resulting from appendicitis. One specimen shows that the duct was embedded in a dense stratum of inflammatory new-formation behind and to the outside of the cæcum, and entirely shut off from the general peritoneal cavity. A perforation had previously taken place. A small foreign body was lodged in a circumscribed cyst cavity. In another case an

autopsy was made upon a patient who had died suddenly from heart disease, and the appendix was found buried in a mass of areolar tissue continuous with a dense band of tissue adherent to a cicatrix upon the skin. This mass of tissue in which was embedded the appendix with a cicatrix denoting a slough was behind the cæcum and the peritoneum. Several years previous this patient had suffered from an attack of appendicitis; perforation had followed and the appendix was placed in a position never again to give rise to any trouble. The condition of perforation was not diagnosed at the time of the attack and the patient never suffered ill consequences from the new position of the appendix. I have also another specimen of an appendix which was situated behind the cæcum and adherent to it. A perforation of it would have been extra-peritoneal, and in that case the danger of a purulent peritonitis would have been avoided. I believe that exceptionally a few of these cases of abscess connected with the appendix are outside of the general peritoneal cavity; but, of course, the large majority of them are intra-peritoneal. Any attempt to excise an appendix during the quiescent period under the circumstances just mentioned would be attended with very great danger, whereas no trouble would ever arise from the condition of the parts.

I know of three cases where the appendix had become adherent to the wall of the cæcum and perforation had occurred through the adherent surfaces directly into the lumen of the cæcum. The escape of pus *per rectum* is the proof offered in these cases of typical appendicitis. The patients recovered and the appendices in their present condition of adhesion with the cæci will probably never give rise to mischief.

To recapitulate, it seems from an examination of the clinical facts and of the pathological material, and from a careful study of the reported cases of appendicitis that excision of the appendix pertains to only eleven per cent. of all the cases of appendicitis. Of these cases represented by eleven per cent. many undergo resolution, some terminate by abscess opening spontaneously or by early incision, while a few are exposed to the dangers of perforation and general peritonitis. Excision of the appendix is suggested as a preventive measure in the small percentage of cases of relapsing appendicitis in which perforation is feared. It has, however, been clearly proved by my specimens that even perforation and general peritonitis are not in all cases fatal complications.

So, then, if those cases of perforation in which recovery follows—and they are more frequent than has been supposed—are deducted from all the cases of perforation, there are left but a few concerning

which the question of the propriety of excision of the appendix would arise. If, now, it is remembered that an incision on the second or third day during the attack in relapsing cases, is attended with most brilliant results, showing a mortality of only a small percentage, is it not wise to observe conservatism and decline to excise the appendix during a quiescent period, when its death-rate is higher than the death-rate of early incision upon the second day? The results which Dr. Willard Parker obtained by operating during the second week after adhesions had formed were brilliant, considering the high death-rate previous to his suggestion. The improvement upon Dr. Parker's operation by early incision upon the second or third day has left but little to be desired in the management of these cases during an attack. Are not, then, a patient's chances for life better after an incision upon the second or third day during an attack than they are after a laparotomy performed during the quiescent period, with the view of removing an appendix which may not be found, and if found cannot be safely removed. Does not such an operation expose the patient to a greater peril than the management of a future attack by early incision?

In conclusion the clinical fact must not be lost sight of that the last attack from which the patient suffered, may have placed his appendix in a position that precludes the possibility of a future attack. From a consideration of these facts it seems wise to condemn the operation of excision of the appendix during the quiescent period as routine practice. Some special reasons should exist to justify the operation—reasons that do *not* exist in the majority of cases of relapsing appendicitis. That occasionally a case may be presented in which excision of the appendix is indicated, I do not deny, but as routine practice in all cases of relapsing appendicitis it is a measure fraught with greater danger than incision and removal upon the second or even the first day of a recurring attack. If this very early incision, as suggested by Sands and practised in several cases by McBurney and others, is resorted to in these relapsing cases, the question of excision of the appendix will arise only in connection with an early attack. If relapsing cases are operated upon at the onset of the trouble the prognosis is better than in those in which later incision is employed. It is one of those operations of expediency which should never be performed, even in a case preëminently suitable, without a thorough examination of the heart, lungs, and kidneys, for under no circumstances, with organic disease, should so bold an operation be resorted to unless the patient is confronted during an attack by death itself. To excise during an interval between recurring attacks in a patient suffering from any organic disease is ex-

posing that patient to a certainty of death, while conservatism might result in a prolongation of human life, during which time no other attack may arise. If an attack should develop, the early operation upon the first or second day in a relapsing case might offer a prospect of recovery that would be more favorable than was at first supposed. At all events, this line of practice does not expose the patient to the danger of death before he is confronted with it, and then it is not too late to expect an escape by a resort to the new and early operation of incision and excision before serious changes have developed that might place the patient's life in jeopardy.

A CLINICAL HISTORY OF TWENTY-TWO CASES OF FILARIA SANGUINIS HOMINIS.

Seen in Charleston, S. C., from 1886 to May, 1890.

BY P. G. DE SAUSSURE, M.D.,
OF CHARLESTON, S. C.

My excuse for this paper is that it deals with a disease scarcely known eighteen years ago, and still but imperfectly understood.

Whether or not filaria have just appeared in Charleston I am not prepared to say, but the case-book of the City Hospital shows that in 1882 or '83 Dr. J. J. Edwards, then a student of medicine, together with several members of the house-staff of the hospital, looked for filaria in the blood of a chyluric patient; and that in 1884 Drs. Edwards and W. D. Bratton examined again and again the blood of a chyluric pregnant woman for filaria, but with negative results. In 1886 Dr. John Guitéras gave the subject a new start, this time with success.

The arduous duties of a large dispensary practice leave me little time and less energy with which to make original research, nevertheless at the end of this article I have set down some facts noted while studying the subject.

CASE I.—A mulatto woman aged about forty. At the time filaria were first found (1886) she was a patient of Dr. Edwards, but had sought treatment of many doctors. She was born in Charleston, S. C., lived in Augusta, Ga., five years, but removed to her birth-place and has lived there ever since. Has had for the last ten years chyluria, with vague lumbar pains, occasional diarrhoea, and during the last three years an irregular fever with chills. In 1884 or 1885 I treated her for lymphangitis of the vessels and glands of the left leg. After some weeks the inflammation seemed to settle in the popliteal space, for in that place I opened an abscess. The pus was not peculiar, but the healing process was slow. During this attack the patient was unable to follow her vocation, that of a seamstress, but before and since she has been able to work; I have lost sight of her for the last two years. This, then, was the first case in which living filaria were found, and

the examination of her blood was suggested by Dr. Guitéras.

CASE II.—A patient of Dr. Edwards. A negro aged nineteen or twenty years. Born and brought up in Charleston, S. C., in the vicinity of which he has spent all of his life. He has spent two or three months of several summers at Mt. Pleasant, a summer resort for Charlestonians, just across the bay. By occupation he was a house-servant. He sought Dr. Edwards's advice for the relief of rapidly developed chylocele which the patient thought was due to traumatism. Dr. Edwards submitted to me some of the fluid drawn from the scrotum and in it I found dead filaria; two nights afterward living filaria were discovered in his blood. Drs. Edwards and Guitéras thought he then had beginning elephantiasis of the scrotum. Shortly after this a mitral regurgitant murmur and dropsy presented themselves. The murmur and ascites increased and the patient died, apparently from heart failure. No post-mortem was allowed.

CASE III.—John G. A nearly pure negro, aged sixty-five, born and brought up in Charleston, S. C., which he claims as his residence, though he has been for months at a time on a rice plantation in South Carolina, at Flat Rock, N. C., and at Newport, R. I. By occupation he is a valet and house-servant. Had perfect health up to 1882, when, following some exertion, he had lumbar pains and hæmato-chyluria, and it was for this that he sought a doctor's advice. In 1883 he came under my care and a careful examination of his urine revealed dead filaria (not then recognized as such), blood, pus and mucous corpuscles, with epithelium from bladder and kidney. A diagnosis of pyuria was made and under tonics and astringents the patient improved, the hæmato-chyluria, lumbar pains and slight diarrhoea returning at intervals, until the fall of 1884, when once more he presented himself for treatment, having then headache, a heart murmur and a swelling of the right side of his neck resembling a unilateral goitre. He again improved under treatment. In 1886, after seeing the filaria in Cases I. and II., I recognized that the objects seen in this patient's urine in 1883 were filaria, so I examined his blood, and found living filaria. In 1889 the patient was once more under treatment for dyspnoea, palpitation of the heart, and persistent headache. In April of that year ascites developed, but under digitalis and iron the patient was relieved. During this attack the patient's heart was embarrassed, but exactly what the trouble was I am unable to make out. The abnormal sounds could not be classified. For the last half year the patient has been able to go about, but any undue exertion distresses his heart and brings on an attack of hæmato-chyluria.

CASE IV.—Mulatto man aged sixty-five, by trade a carpenter, born in Charleston, S. C., which place has been his home, but he, like most of the patients reported, has from time to time lived in other parts of the State. His health was perfect up to seven years ago, when he had his first attack of hæmato-chyluria, which was preceded by lumbar pains, abdominal colic, and dysentery. His medical attendant diagnosed cholera morbus. In 1887, when he

came under my care, his blood was examined and filaria found.

CASE V.—Allen, a negress, aged forty-two, born in South Carolina. Has lived in Charleston for the last twenty years, but is not clear as to her previous residence. By occupation a cook. Has suffered from chyluria for five years. No other trouble until three years ago, when she had back-ache, colic and slight dysentery, but did not stop working. Filaria found in the blood in 1886. She is a widow, has had several children, only one of whom is living. She does not know of what her husband or children died. Looking for some hereditary influence, I asked that her child should come to the office.

CASE VI.—Allen's son, a lad aged eighteen years, born and brought up in Charleston. Is an epileptic, but otherwise is in excellent health. No pains nor chyluria. Blood examined in 1887 and filaria found. The filaria do not seem to give this patient any trouble. Bromides control his convulsions more or less.

CASE VII.—Mr. B., thirty-three years old, born in Charleston. Is now a constable, but has followed different occupations, one of which carried him for several months to Nassau, where he had "fever" during the late war. In 1870 he had his first attack of chyluria. Between 1870 and 1880 he had a peculiar inflammation of the glands of the groin. He says his physician could do nothing with them, and that finally an abscess formed. Since 1887 with his chyluria he has lumbar pains and some dysentery. Filaria were found in 1887. He does not know if they had previously been looked for.

I think this patient is mistaken in regard to his age, which I would say is about forty-three.

CASE VIII.—Mr. Y., aged about twenty-seven, born in Charleston, S. C., where he has spent all of his life. On account of a diseased hip-joint he has never been able to do much manual labor, though he has been a great deal about the shipping. Excepting hip disease he enjoyed perfect health until four years ago, when, following a liberal meal at a picnic, he was seized with cholera morbus. From then until now he has been subject to attacks of dysentery with abdominal pains. Examined for and found filaria in 1888.

CASE IX.—White female, aged thirty-five; born and raised in Charleston, though she has visited for two or three months on various plantations in the vicinity of Charleston. She is married, and at the time of first observation in 1887 had one child. Her health was good until the latter part of her first pregnancy, when she began to suffer with chyluria and severe pains in the lumbar region. The urine was white until after labor, when it became natural and remained so until the eighth month of her second pregnancy, when the urine became white again. I saw her for the first time at this period and found chyluria and rectal tenesmus. The urine contained vesicular masses large and small, the former about one and one-half inches long by one inch broad. These vesicles were entirely new to me and I can only describe them as looking like egg albumin enclosed in a fine membrane, the whole being

in color and transparency like uncoagulated albumin. Sometimes they would be arrested in the urethra and pass only after long bearing-down efforts. I examined several of the masses, for they could be handled, though soft and compressible, but I could make nothing out of them. Once I thought I had a Bilharzia ovum, but of this I am not sure. I can only say that the vesicles and what I saw in one of them under the microscope were new to me. In this patient's blood filaria were found a few hours before labor, as well as the night before. Immediately after labor none were found either in blood from the cord or in that drawn from her finger. Examination of the blood and milk on the following day was also negative and no filaria were found in the child's blood. After some weeks chyluria disappeared, but filaria were again found in the mother's blood (child's blood was always negative). Patient became disgusted with so many examinations and I lost sight of her until 1890, when once more she consulted me and I found that she had been confined again in 1889. She said that she had had no trouble during pregnancy, but this I doubt, and think she made the statement to keep my needle out of her baby, to whom I had been summoned to relieve it of dysentery. The whole family—father, mother, and three children—were examined, and I got negative results and my discharge. The child has since died under the care of another physician. The certificate of death shows enterocolitis as the cause.

This case is particularly interesting, for the discharges from the four-months-old infant were identical in appearance with those of the other patients suffering from chyluria and dysentery.

CASE X.—Mr. G., aged fifty-seven, born in Ireland, has lived in Charleston for the last 35 years. He has been a wanderer and is a "jack of all trades." Since 1875 he has been much of the time in hospital, complaining of vague pains. In 1887 he applied for relief of lymph scrotum and enlarged groin glands. Filaria found in blood glands, which were removed, but were not adult filaria.

CASE XI.—S., aged fifty-eight, born and brought up in Charleston. Present occupation that of a house-cleaner, but like most negroes, has turned his hand to everything, and I must say of him as I have of others, that he claims Charleston as his residence, but acknowledges visits to other places; for instance, he was in Savannah, Ga., for four months. He sought relief in 1888 for chyluria of several months' standing, with lumbar pains and dysentery. Filaria found, and from eye measurement I would say that they were larger than the average. This patient had, during the month preceding his coming to me, an irregular fever with chills; no enlarged glands present.

CASE XII.—J., aged fifty-five, born in South Carolina. Resided for last ten years in Charleston; before that in different parts of the State, with visits of months' duration to North Carolina and Georgia. His work at times has been loading and unloading vessels. Has had chylocele for four years. This

rapidly growing larger, he applied for treatment in 1889. Blood examined and filaria found. On being told that an operation would have to be performed he left, and I have no further history.

CASE XIII. (Should have been reported amongst cases found in 1886.)—A negress, one hundred years old, born in Africa. Was brought over to South Carolina in a slave-ship.

With the exception of four years, she has lived in Charleston. Called to attend her in 1886 for a cold abscess, pointing just over the scapula. Her age prevented her from giving much of a history. Thinking that her birth-place gave her a right to filaria, I examined her blood and found them. She was sent to the hospital, where the abscess was opened, but nothing peculiar was noted about the pus. She has since become insane, and has been for the last three years in an asylum. The assistant physician writes, May 10th, that "she has occasional chyluria, but no glandular swellings, and, in fact, is remarkably well." Filaria were found two weeks previously.

CASE XIV.—Mr. J., aged forty-five, patient of Dr. J. L. Dawson, Jr., who gives the following history: Birth-place and residence Charleston, S. C., but has lived in Florida, Georgia, and North Carolina. Has had chyluria for years. An attack is always preceded by lumbar pains. Is married and the father of several children, all of whom are healthy. This patient's occupation for the last few years has been that of a book-keeper to a large fruit firm, and from time to time he drinks water on vessels from Cuba and the West Indies. Had chyluria before this, however. Filaria found in 1888.

CASE XV.—Mr. M., aged thirty-two, lawyer, a patient of Dr. Kinloch's; birth-place and residence Charleston, S. C. Filaria found in blood by Dr. Kinloch in 1887; subsequently in urine and blood by Dr. Guitéras. Chyluria is the only symptom that he presents.

CASE XVI.—Mrs. B., aged fifty-two; born in Germany, but has lived in Charleston, where she came direct from Germany thirty years ago. By occupation a cook. Has had chyluria for the last eight years, but no other symptoms, save lumbar pains. Filaria found in 1887. Has one child, a boy, born since her arrival in this country. The boy's blood gives negative results.

CASE XVII.—M., male, aged about sixty years; born in South Carolina. Lives now in Charleston, but has lived in different parts of the State. In good health until several months ago, when he began to have trouble with his urine, which he describes as muddy and streaked with blood. Applied for treatment of cystitis. Dr. Edwards thought that the disease might be caused by stone, but the sound gave negative results, so we determined to look for filaria, which were found in his blood in 1890. This patient complains of great muscular weakness.

CASE XVIII.—Mrs. F., aged twenty-six, a patient of Dr. Manning Simons. Born and raised in Charleston. Is married and has two children. Has had chyluria for fourteen years. Has also had irregular fever with chills, and enlargement of the lymphatics

of left leg and arm. None of these have ever suppurated, and the patient is able to go about attending to all of her household duties. Filaria found in 1890.

CASE XIX.—Sallie, aged forty-five, patient of Dr. D. Ryans. Born in South Carolina, lives in Charleston now, but gives no clear history of her birth-place or residence before she came to Charleston. Gives no account of the duration of chyluria, which is her only trouble. Is married and the mother of several children. Filaria found in 1890.

CASE XX.—Frank, aged forty-five, patient of Dr. Ryans. No history, save that he was born in South Carolina and has lived in Charleston for many years. Applied for relief from chyluria, which he says he has had for years. Filaria found in 1890.

CASE XXI.—Martha R., aged forty-eight; born in South Carolina; residence for last twenty-six years, Charleston. Health good until seven months ago, when she had an attack of hæmaturia, consequent upon gravel, her physician thought. Since then she has suffered from palpitation of the heart and dyspnoea. Three weeks ago oedema of the lower extremities appeared. On May 13, 1890, presented herself for treatment. Filaria found.

CASE XXII.—Gibbes, aged sixty, born in South Carolina; residence for last thirty-six years, Charleston. Health perfect until six months ago, when he had an attack of palpitation of the heart with dyspnoea; this has grown rapidly worse for the last month. Oedema of feet and lower extremities now. Filaria found, May, 1890.

Tabulating the cases, I find that the disease was distributed as follows: 9 colored males, 6 colored females; 5 white males, and 2 white females. Of the 22 cases 16 were in married people, all of whom had children. In over one-half of the cases filaria were searched for in children of the married, in husbands or wives, or in friends and bed-fellows of the single ones; and yet but in a single instance did I find two cases in the same house (Cases V. and VI.). In the child of Case IX. I think, if the search could have been carefully carried on, filaria would have been found.

As to the social conditions of the patients, 15 were colored, 7 white; of the latter only three can be said to be in high life. All of the others, black and white, belong to the lower and middle classes. Why this difference? Is it from abode or diet? The houses of our domestics and laborers are very similar to those of the same classes elsewhere.

In reference to the diet and water, the former is that of the working-class in other cities, only that here they eat a great deal of fresh fish—in fact, during the summer months fish of some sort takes the place of meat. As to the drinking-water, I differ absolutely from Dr. Guitéras, for he considers it mainly cistern, and I know from a ten-years' practice amongst the lower classes that it is mainly

pump or artesian, and exceptionally cistern water. Sometimes, for a short period just after rain-storms, the poor have barrel water, but the supply soon runs out, and they go back to the pumps and artesian wells. A word about the barrel water, which is rain-water caught in barrels and casks open to dust and dirt, and one can always find with the microscope many living things in the water. The same is true of the artesian water, which, though it comes from wells 1800 to 1900 feet deep, is, before being served, stored in an open reservoir, which, to the best of my knowledge and belief, is bricked half-way down the sides, but only puddled at the bottom. I am not enough of a microscopist to say to what class the animals found belong, but know this, that with the microscope I have spent hours seeing them fight and destroy each other. The better class of Charlestonians drink cistern water. As to mosquitoes, I think they are as common in the homes of the rich as in those of the poor.

Diagnosis of the disease can alone be made with the microscope, for by reference to the table attached to this paper we find that chylocele, chyluria, lymph-legs, and glandular enlargements are frequently found, yet examinations of the blood may fail to show filaria. Again, we find filaria without any of these disturbances (Cases VI., VII., X., XXI., and XXII.).

There is no pathognomonic symptom. Sometimes we have fever and chills, at other times neither. The disease, as a rule, does not appear to prevent the patients from working. Cases II. and III. are exceptions, and some of the other patients were for a time disabled, but returned to their duties, and the filaria were still found when looked for. The chyluria is intermittent, and in no case that I have seen was it always present; and I would state further, that during its presence a given specimen of blood would not show more filaria than another specimen examined when chyluria was absent. In but two cases have I ever failed to find filaria after having once seen them (Cases VIII. and IX.). In Case III. his residence for three months at Flat Rock, N. C. (at an elevation of 2000 feet above the sea-level), always reduced the number of filaria, but after being in Charleston for eight months they became numerous again. Would they entirely disappear if he remained in the mountains?

To show the impossibility of making a diagnosis without the microscope, I would state that during the last three years I have examined three cases of chylocele with negative results (blood of two, fluid of one). Six cases of chyluria and one of hæmato-chyluria have also given negative results, though one of these may yet prove to be a case of filaria, for it occurs in a patient who is a printer, and as yet I have not been able to examine him at the end

of his work, which is about 5 A.M. Five cases of elephantiasis of the leg have been under observation, but no filaria were found.

Learning from our veterinary surgeon, Dr. Benjamin McGinnis, Jr., that filaria were very common in the dogs of Charleston, with his and Mr. L. W. Bacaise's assistance I made some observations on the disease in the dog, and I found that the embryo filaria in man and dog are identical—those of the man being possibly a little larger and their movement more active and strong. Very early in the investigation I was surprised to find that in the dog the embryo filaria could be found at any time of the day or night. In man it is different for I have been able to find them in the day but twice, once at 10 A.M. in the negress from Africa (Case XIII.), and once at 5 P.M. (summer) in John G. (Case III.)

Carrying the investigation still further, I examined fleas from filarial dogs, but got negative results until the morning of May 14th, when I found two filaria in the body of a flea which was taken from a diseased dog. This observation will, I hope, be confirmed, for it may throw considerable light on the mode of infection. I had purposed to make some examinations on the puppies of an infected bitch, but time has been wanting. I therefore would suggest that anyone interested in the subject may, by such examinations, clear up the question of heredity.

In the dog the disease causes death from ascites, which seems to be in consequence of the adult filaria collecting in the heart, a condition that we found in five post-mortems.

It is worthy of remark that in the one case of death occurring in a patient known to be afflicted with filaria, the cause of death was "dropsy with heart failure"—no post-mortem was allowed, so the presence of the adult filaria in the heart was not proven. It is suspicious that Cases II., III., XXI., XXII., should have heart murmur and dropsy.¹

A REPORT OF TWO CASES OF STRICTURE OF THE URETHRA AND CHRONIC PROSTATITIS PRESENTING INTERESTING REFLEX SYMPTOMS.²

BY THOMAS R. NEILSON, M.D.,

INSTRUCTOR IN GENITO-URINARY DISEASES IN THE UNIVERSITY OF PENNSYLVANIA, SURGEON TO THE EPISCOPAL HOSPITAL AND TO ST. CHRISTOPHER'S HOSPITAL FOR CHILDREN, ETC., PHILADELPHIA.

THE two cases which I have the pleasure of reporting in this paper have seemed to me to possess sufficient interest to warrant their being recorded,

¹ I would beg to return to Dr. Louis R. Staudenmayer, of Charleston, my thanks for his help in making microscopic examinations and preparing this paper.

² Read before the Philadelphia County Medical Society, May 28, 1890.

because of the marked and peculiar symptoms associated with the disease of the urinary outlet present in each.

CASE I.—Mr. —, a well-developed, athletic young gentleman, twenty-four years of age, came to my office in December last, giving me the following history: About a year previously he had had what seems to have been an attack of subacute urethritis, following an exposure. For this he used, by advice, a strong solution of sulphate of zinc as an injection, with the result of aggravating the symptoms to a considerable degree. Later in the attack a catheter, which entered the canal with great difficulty, was passed, and again the symptoms were for a time increased. Gradually the discharge and other symptoms subsided, but never entirely disappeared. At the time of his first visit to me he complained of frequently recurring attacks of scalding on micturition, accompanied with a purulent or muco-purulent discharge, worse after defecation, and with an annoying itching just within the meatus. At times this itching was felt over the glans penis, the scrotum, and in both groins. In addition, a sharp pain, neuralgic in character, was felt just below the umbilicus, coming at irregular intervals, lasting for a few moments, and then suddenly disappearing. Simultaneously with the occurrence of this pain, a tickling sensation in the throat was felt, and a short, laryngeal cough was provoked.

The patient had observed that friction over the umbilical region, such as that occasioned by using a towel after bathing, would aggravate the itching in the anterior portion of the urethra when it was present, or produce it when it was not noticed. In addition to the urethral trouble, the patient suffered slightly from internal hæmorrhoids, and when these were congested, during constipation, the itching in the urethra was slightly increased.

Upon examination, I found the circumference of the penis to be three and three-quarters inches, the meatus very small, only 12 (F.) calibre, and the anterior portion of the urethra excessively sensitive to the small-sized rubber bougie which could be introduced into the orifice.

The first step in the treatment consisted in the division of the meatus, so that it would easily admit a No. 34 sound. Being then able to further explore the urethra, I discovered a stricture of 26 calibre at two and three-quarter inches from the meatus, and a gleety condition in the bulbous portion. The whole canal, especially the posterior portion, was extremely tender, and a characteristic prostatic discharge was brought out on the shoulder of the bulbous explorer. Subsequent treatment consisted in passing at regular intervals full-sized steel sounds, the use of hot sitz baths, and the local employment of a sedative and astringent lotion for the itching of the scrotum and groins. After the sounds had been passed a few times the tenderness of the urethra disappeared, micturition was of course much improved (owing to the enlargement of the constricted meatus), the itching sensation in the urethra was no longer observed, and the pain near the umbilicus and the tickling in the throat ceased. It is inter-

esting to note that at first, during the employment of the sounds, the latter symptoms were always aggravated for a short time after each visit.

By the first of April the patient considered himself well, but a little later, after taking a heavy cold and being indiscreet in his diet, the prostatic trouble was excited again, being manifested by a mucous discharge from the urethra while at stool, and a return of the itching near the meatus, but less intense than formerly. Under treatment this condition has again subsided, and the patient is now practically well.

CASE II.—Mr. —, also a private patient, a gentleman twenty-three years of age, having the history of an attack of gonorrhœa, complicated by prostatic cystitis, three years before he first consulted me, which was at the beginning of this year. The symptoms present were a slight gleety discharge, and burning pain on micturition and occasionally during the intervals. Whenever this pain was present there was also felt a similar pain on the inner border of the left foot near the first metatarso-phalangeal joint, and extending to the ball of the great toe. Friction of the glans penis by contact with the clothing produced the painful sensation, both in the urethra and in the foot; and, conversely, rubbing the foot upon a carpet or rug, or warming it before a fire, would excite the sensation in the urethra.

Examination showed: Circumference of penis three and a half inches, a stricture of 24 calibre at two and a half inches, and a highly sensitive condition of the urethra, most marked in the posterior portion, from which the discharge of chronic prostatitis was obtained by the explorer. I advised internal urethrotomy for the stricture, but my patient would not consent, preferring to trust to the less certain method of gradual dilatation. Prescribing a mixture of the bromide of potassium, tincture of hyoscyamus, and camphor water, for the purpose of reducing the abnormal sensitiveness of the urethra, I began the treatment with sounds, and, as the condition of the urethra improved, the painful sensations, both in that canal and in the foot, became less frequent and less intense, and had nearly disappeared when an attack of illness necessitated the discontinuance of the treatment. After recovering from this attack my patient left the city, by the advice of his physician, and on his return, recently, tells me that he again has his old sensations, and has decided to submit to the operation of internal urethrotomy.

REMARKS.—The existence of pain or other morbid sensation in association with or arising from stricture of the urethra, cannot, of course, be considered exceptional. All authorities upon the subject of stricture mention among its symptoms (exclusive of those at the seat of the lesion or lesions) pain in the perineum, the hypogastrium (when cystitis is present), the small of the back, the loins, the testes and groins, and sometimes extending down the thighs. Sensory disturbances at more remote situations are certainly exceptional.

A case similar to the second here reported was described by Sir Benjamin Brodie, in his "Lectures Illustrative of Certain Nervous Affections" (*Dunglison's American Medical Library*: "Medical and Surgical Monographs," Phila., 1838, p. 38). The case was that of a gentleman who consulted him on account of a pain in the instep, so severe as to make him lame. Incidentally, the existence of a stricture was discovered, and its dilatation was followed by the relief of the pain.

I have not been able to find any reference to a case resembling the first one of this paper, certainly not so far as the throat symptoms are concerned.

MEDICAL PROGRESS.

Quinine and Sulphur in the Treatment of Diphtheria.—According to the *Provincial Medical Journal*, DR. BURG-HARDT employs equal parts of quinine and flowers of sulphur in the local treatment of diphtheria. The powder is blown upon the false membrane, after which, if the disease is in the pharynx, the patient is not permitted to swallow anything for an hour and a half. As a prophylactic measure the powder is also insufflated into the nasal cavities, even when these parts are not infected. Of the thirty three cases treated in this manner by Burghardt all recovered, and in none was the powder applied oftener than twice daily. Internally he prescribed the tincture of the chloride of iron. In addition, the patients were bathed with vinegar, and were given wine, brandy, and milk.

Therapeutics of Green Coffee.—DR. LAUDARABILCO recommends an infusion of green coffee in the treatment of gout, gravel, nephritic colic, and migraine (*Journal de Médecine de Paris*). The following varieties of coffee are used: Martinique, one-half; Mocha and Bourbon, of each, one-fourth. At night six drachms of the mixture are placed in a glass of water and macerated about twelve hours. The contents are then stirred and strained, and the clear liquid drunk without sugar, and while the stomach is empty. Food may be taken soon after. The results are said to be extremely good.—*Provincial Medical Journal*, June, 1890.

Thallin in Typhoid Fever.—DR. FRANZ SCHMID has published an inaugural dissertation on the treatment of typhoid fever with thallin, his observations being made upon twenty-five cases. The drug was administered in doses varying from $\frac{1}{4}$ grain to 3 grains every hour. The author draws the following conclusions:

1. In the above-mentioned doses thallin has an excellent antipyretic action in typhoid fever of moderate severity. In severe cases it is not very effective. In the former cases it has a good effect on cerebral symptoms.

2. The drug does not cause collapse nor renal irritation, and it is without disagreeable effects on the heart and lungs.

3. It seems to reduce the mortality of typhoid fever, but has no appreciable effect upon the duration of the disease.

4. The thallin treatment is of as much value as the hydropathic treatment, and is, of course, less annoying to the patient.

In the *Correspondenzblatt für Schweizer Aerzte*, No. 9, 1890, Ruetimeyer writes that he has never seen collapse following the use of thallin, even when given in much larger doses than those of Schmid, and that even in severe forms of the disease it has a good effect on cerebral symptoms.—*Provincial Medical Journal*, June 2, 1890.

Prescription for Eczema in Children.—The following prescriptions for eczema in children are quoted in the *Annals of Gynecology and Paediatrics*:

1. R.—Bismuth subnitrate . . . 4 drachms.
Zinc oxide . . . 1 drachm.
Carbolic acid . . . $\frac{1}{2}$ drachm.
White vaseline . . . 2 ounces.—M.
2. R.—Bismuth subnitrate . . . 2 drachms.
Zinc oxide . . . $\frac{1}{2}$ drachm.
Glycerin . . . 1 $\frac{1}{2}$ drachm.
Carbolic acid . . . 20 minims.
White vaseline . . . 26 drachms.—M.

The latter is a very elegant and soothing ointment.

If there is much stinging and irritation the following lotion will be found valuable:

- R.—Bismuth subnitrate . . . 1 drachm.
Glycerin . . . 4 drachms.
Carbolic acid . . . 12 minims.
Rose-water to make . . . 1 ounce.—M.

Shake, and apply with a soft brush.

Cancer of the Pancreas; Extirpation; Recovery.—A successful extirpation of a primary cancer of the pancreas is reported by PROFESSOR RUGGE, of Bologna (*Medicinische-chirurgische Rundschau*, April 15, 1890). The patient, a woman of fifty years, came to Rugge's clinic in August, 1889, complaining of constant gastric pain, anorexia, and obstinate constipation. The pain began in the epigastrium, spreading from there to other parts of the abdomen. Physical examination revealed a tumor in the neighborhood of the transverse colon in the left hypochondrium and surrounding the umbilicus. The tumor was rather irregular in shape, felt firm on pressure, and measured about ten inches in length and four inches in width. The author made the diagnosis of probable retro-peritoneal sarcoma.

The patient insisting on the removal of the tumor, operation was undertaken on September 4th. The removal was rendered extremely difficult by the softness and fragility of the growth and numerous adhesions. Finally, extirpation was accomplished and the growth was found to be the pathologically changed pancreas, which the microscope proved was carcinomatous.

Healing of the wound was most rapid, and the general condition of the patient improved from day to day, and notwithstanding the fact that she desired and received a mixed diet, digestive disturbances diminished and finally disappeared. The author thinks that this remarkable case shows that after total extirpation of the pancreas other glands perform its functions, particularly Brunner's glands and Lieberkühn's follicles.

THE MEDICAL NEWS.

A WEEKLY JOURNAL OF MEDICAL SCIENCE.

COMMUNICATIONS are invited from all parts of the world. Original articles contributed exclusively to THE MEDICAL NEWS will be liberally paid for upon publication. When necessary to elucidate the text, illustrations will be furnished without cost to the author.

Address the Editor: H. A. HARE, M.D.,
1004 WALNUT STREET,
PHILADELPHIA.

Subscription Price, including Postage.

PER ANNUM, IN ADVANCE \$4.00.
SINGLE COPIES 10 CENTS.

Subscriptions may begin at any date. The safest mode of remittance is by bank check or postal money order, drawn to the order of the undersigned. When neither is accessible, remittances may be made, at the risk of the publishers, by forwarding in registered letters.

Address, LEA BROTHERS & CO.,
Nos. 706 & 708 Sansom Street,
PHILADELPHIA.

SATURDAY, JUNE 28, 1890.

CODEIN.

ACCORDING TO LOEWENMEYER (*Deutsche medizinische Wochenschrift*, No. 20, 1890), codein is very little employed in Germany, notwithstanding the encomiums which it receives from physicians elsewhere and especially from the French.

He therefore recently instituted an extensive series of observations in Jacobson's service at the Jewish Hospital in Berlin. Some five thousand doses of the drug were given to about four hundred patients, some of whom took it for weeks, and others for months; yet in no case did he observe any untoward results. He therefore recommends its use in place of morphine, rating it as a narcotic of somewhat lesser intensity than the latter, but on the whole superior, because of its freedom from danger. He especially endorses Dr. Lauder Brunton's assertion of the peculiar applicability of codein to the relief of painful affections of the abdomen. Gastralgia, colic, and the various visceral neuralgias seemed peculiarly amenable to its influence. The patients had long periods of respite from pain. They found sleep, and when, after a certain time, new exacerbations occurred, the medicament in the same dose as at first still exerted the same favorable influence.

Far more important than the relief of functional disturbance, however, in which the psychic factor

may be presumed to have more or less influence, is the possibility of relieving the pain dependent upon organic disease, gastric cancer, cancer of the liver, cancer of the intestine, etc. While morphine was effective in ninety out of one hundred cases of this nature, codein succeeded in only sixty.

The question as to the indications for one or the other of these drugs is quite difficult to answer. Excluding idiosyncrasies, Loewenmeyer is disposed to think that attacks of pain of great intensity, occurring in paroxysms, are not likely to be favorably influenced by codein, but are alleviated by morphine. This was particularly noticeable in hepatic colic and in renal colic. He would also consider it a mistake to administer codein in cases of gout or of circumscribed inflammation of the peritoneum, in which large doses of opium act so happily in preventing extension of the inflammatory process. After the acute symptoms have subsided, however, codein is useful in controlling the remaining pains.

Incomparably more favorable than in any affection heretofore mentioned, is the action of codein in diseases of the thoracic viscera, and particularly of the organs of respiration. This was especially notable in cases of phthisis, alike for its influence upon the cough, for relief of pain, and for modifying acute bronchitic or pneumonitic exacerbations. Good results were likewise evident in some cases of bronchial asthma. The drug was also administered in some cases of severe cardiac lesions in which morphine had previously been employed, and without seeming to produce any toxic effect. On the other hand, it was of little avail either in organic or functional nervous affections. As a hypnotic, however, it acted readily, nor did it seem to have a tendency to produce an unfortunate habit, such as follows the continued use of morphine.

While Loewenmeyer cannot agree with the opinion that codein is useful in the direct treatment of the morphine habit, he looks upon it as a prophylactic, inasmuch as by its substitution for morphine the contraction of the habit may be avoided.

We have called attention to this careful study of an important drug, because we believe that in America, too, its use might be profitably increased. Our experience tallies closely with that of Loewenmeyer so far as reported—an important omission in the report being its use in diabetes mellitus—and the concluding observation of that author is one to which great heed should be given. Many unfortunate cases of morphine habit have been due to un-

guarded prescription of the drug, or too long continued resort to it, by physicians. It is much easier to avert contraction of the habit than to cure it. Therefore, if codein be applicable at all in cases where an opium alkaloid is indicated for anything more than occasional use, it should be preferred. We are inclined to go still further than Loewenmeyer and to urge caution and reluctance in the use even of codein for prolonged periods. Let continuous resort to any drug of this nature be only adopted after careful deliberation, as an absolute necessity, and with the safeguard, to both patient and physician, of a consultation.

CORRESPONDENCE.

FIVE CASES OF PTOMAIN POISONING.

To the Editor of THE MEDICAL NEWS,

SIR: I take the liberty of presenting these cases, which to me were of considerable interest, to my colleagues in the medical profession for their consideration. I think that the report may perhaps assist us in finding the etiology of some of those most serious and painful conditions which we so often meet with, and where the cause is somewhat obscure.

I had been practising for some time in the mining regions near Charleston, S. C., a dangerously malarial section, and while there had seen some of the most desperate and malignant forms of malarial affections, such as malignant malarial dysenteries, paralyses, and congestions, and so was not in a condition to be surprised by anything I met with. However, on Monday, in the latter part of April, 1890, I met with a series of five cases which specially attracted my attention after the death of two of them. On the previous Friday evening there had been a beef and a hog killed, the meat of which was sold in the local market on Saturday; it had not been kept on ice, but the weather was pleasant, not hot. My patients ate some of this meat Saturday afternoon and again on Sunday morning—four women and one man. One of the women at about eleven o'clock Sunday morning was taken with apparently a violent attack of cholera morbus, in which condition she remained until I saw her early on Monday morning. Another of the women was attacked in the same way at about 1 o'clock Monday afternoon; she had eaten some of the pork. The other three were attacked in a similar way during Monday.

The meat had appeared to be perfectly fresh and very nice, so that when I inquired as to whether they had eaten anything which might have caused the trouble, they did not mention the meat. The first two women I found in the following condition: Lying with the knees drawn up and suffering great pain, paroxysmal in character, abdomen distended, tympanitic and painful on pressure, countenance anxious, great nervous excitement, temperature 101° , pulse rapid, bowels constipated, no history of a chill or any previous debility. I considered this condition as cholera morbus, although

there had been but one slight attack of vomiting and no purging, so I gave a large enema of warm water and soap, thinking that I would clear the bowels of the irritating substance. The enema acted well and brought away large quantities of fecal matter. I then gave a hypodermic injection of one-half grain of morphine, ordered hot stupes to be applied over the abdomen constantly, and one-quarter grain of morphine to be given every three hours until relieved.

On calling next morning I found that my patients had had little or no relief, and that a dysenteric discharge had set in, temperature 102° , pulse rapid and compressible. Thinking this to be dysentery following cholera morbus I gave ipecacuanha, opium, and bismuth-subnitrate; also quinine, five grains, three times a day. But the condition continued to grow worse and by night my patients were in collapse, and died the next morning (Wednesday).

Of the other three, the man recovered on the above-mentioned treatment, although he went through a troublesome and severe attack of dysentery. On Tuesday it was suggested to me that perhaps the meat was the cause of the trouble, and the idea struck me as being a likely one and suggested ptomaines. I immediately put the other two women on the following treatment: To one, after washing out the lower bowel with solution of bichloride of mercury 1:3000, I gave bichloride of mercury one-sixteenth grain every four hours; she recovered rapidly. To the other I gave tincture of chloride of iron, 20 drops every four hours, also an enema of the bichloride of mercury solution; she recovered, but slowly.

On examining into the histories of these cases I found that all of them had eaten either the pork or the beef. The man's case was milder than the others, which fact, I think, was due to the beef eaten by him having been parboiled before being roasted, which he states was the case. I have also been informed that numbers of others had dysentery after eating meat, on several occasions previous to this. Finding that a considerable amount of distemper existed among the hogs, cows, and horses, I concluded that perhaps, the animals being in an unhealthy condition, putrefactive changes set in rapidly, and, although being only incipient—for the meat seemed fresh—ptomaines were formed, and that my patients were suffering from poisoning from this product.

Now in the section where I have been practising for a year fresh meat could only be had once a week—on Saturday—and I found, on looking over my cases of dysentery for the year—and a great number of them there were—that most of them were in the spring and summer, and that frequently fresh meat had been eaten a day or so before, and that when such was the case the onset of the disease was similar to that of the above-mentioned cases, varying in severity, due sometimes to the mode of cooking and again to the quantity eaten; although there were not many deaths, I must confess my treatment of that disease with ipecacuanha, etc., was not satisfactory.

The theory I wish to endorse is that as dysentery, cholera morbus, cholera infantum, etc., are most common in the summer months, in hot and filthy quarters, and in the tropics, and that as the summer months and warm climates are the conditions most apt to convert albuminoid substances into ptomaines, may not ptomaines,

more frequently than we generally suppose, be the cause of these affections? And that, if such be the case, could not a more satisfactory treatment than the present be formulated? There was a condition in my cases which I failed to make much mention of above, which, I think, goes to strengthen the ptomaine theory. It was the great nervous excitement, which appeared to me to be due to the absorption into the blood of a virulent septic poison; for the local intestinal conditions I did not consider severe enough to cause it; also the collapse I considered due to this cause.

I did not make an autopsy of either of these bodies or examine the discharges chemically, but I feel convinced that they were cases of ptomaine poisoning, and that the above-mentioned diseases are frequently caused by this product. W. MAZYCK MEMMINGER, M.D.

CHARLESTON, S. C.

NEWS ITEMS.

THE next meeting of the Fifth District Branch of the New York State Medical Association will be the eighth special meeting, to be held in Kingston, Ulster county, New York, on Tuesday, July 22, 1890. A large attendance is desired.

THE 17th annual session of the Mississippi Valley Medical Association will be held at Louisville, Kentucky, October 8, 9, and 10, 1890. The medical profession is respectfully invited to attend. The meeting promises to be of great social and scientific interest, as the profession of Louisville are doing their utmost to make it a success. Ladies accompanying physicians will be made especially welcome. Gentlemen wishing to read papers will please send titles as soon as possible to the secretary, Dr. E. S. McKee, 57 W. Seventh street, Cincinnati. The American Rhinological Association will also meet in Louisville October 6, 7, and 8, 1890.

A Medical Bishop.—The Right Reverend Henry Callaway, Bishop of Saint John's, Caffraria, South Africa, who died March 26th, was a regularly qualified practitioner of medicine and surgery, his M.D. degree having been granted by the King's College, Aberdeen, in 1853. He was in his seventy-third year at the time of his decease.

Dr. Virchow has recently been on a brief visit to Hisarlik, where his old friend Schliemann is still engaged in explorations. On his return, while passing through Constantinople, he was made the recipient of marked attentions by the Medical Society of that city. He addressed the Society in response to remarks by the President, Dr. Stekulis, saying that there was a manifest and happy awakening in many parts of the Orient, of which he himself had been a witness, and the scientific world was destined to be the gainer by the contributions of countries that had hitherto been receivers only, and giving out nothing of value.

Professor Nussbaum Pensioned.—Dr. von Nussbaum, it is announced, will be awarded a pension by his government in consideration of his distinguished services in surgery. He has been invalidated on account of a severe attack of influenza, about six months ago, which has left him partially paralyzed and nearly blind.

THE University of Halle now has its Hygienic Institute, with suitable lecture-hall and laboratories. Professor Renk has been appointed to be its director.

Obituary.—The death of Dr. Francesco Scalzi, at Rome, is the subject of memorial mention in the *British Medical Journal*. Professor Scalzi was the most eminent authority in materia medica, in Italy, of his generation. He established, in 1876, the first and the best, in Italy, Institute for Experimental Pharmacology and Toxicology. He was a distinguished lecturer and teacher, an editor, the writer of 112 publications on various subjects, a hospital physician, a sanitarian, and a public official. He was fluent, eloquent, fertile, and withal blessed with a sound judgment. He was in his 70th year.

Large Russian Surgical Fees.—Dr. Sklifissowsky, of Moscow, having been called to Odessa to operate upon a case of hip-disease, was presented with an honorarium of 11,000 roubles, or the value of £1,222. This is, for Russia, an enormous fee, and the surgeon, with remarkable generosity, passed it over untouched to the University of Odessa, to found a medical scholarship. He also was called upon, incidentally to his other consultations, to operate upon a cancer of the breast, for which he received 2000 roubles. These large fees are for the elect only, the average Russian medical man being very poorly paid.

OFFICIAL LIST OF CHANGES IN THE STATIONS AND DUTIES OF OFFICERS SERVING IN THE MEDICAL DEPARTMENT, U. S. ARMY, FROM JUNE 17 TO JUNE 23, 1890.

By direction of the Secretary of War, leave of absence for six months, to take effect when, in the opinion of his department commander, his services can be spared, is granted WILLIAM O. OWEN, JR., *Captain and Assistant Surgeon*.—Par. 12, S. O. 139, A. G. O., June 14, 1890.

By direction of the Secretary of War, EUGENE L. SWIFT, *First Lieutenant and Assistant Surgeon*, is relieved from duty at Fort Spokane, Washington, and will report in person to the commanding officer at Fort McDowell, Arizona, for duty at that station, relieving Marlborough C. Wyeth, *Captain and Assistant Surgeon*, who, on being thus relieved, will proceed to Fort McIntosh, Texas, and report in person to the commanding officer thereof for duty at that station. The officers named will also report by letter, upon their arrival at their new stations, to their respective department commanders.—Par. 10, S. O. 140, A. G. O., June 16, 1890.

By direction of the Secretary of War, leave of absence for two months, to take effect July 1, 1890, is granted CHARLES B. BYRNE, *Captain and Assistant Surgeon*.—Par. 13, S. O. 141, Headquarters of the Army, A. G. O., June 17, 1890.

By direction of the Secretary of War, leave of absence for four months, to take effect upon the final abandonment of Fort Maginnis, Mont., is granted to WILLIAM W. GRAY, *Captain and Assistant Surgeon*.—Par. 14, S. O. 141, Headquarters of the Army, A. G. O., June 17, 1890.

By direction of the Secretary of War, SAMUEL M. HORTON, *Major and Surgeon*, will visit the encampment of the Maine Volunteer Militia, at Augusta, Maine, during the period of its encampment, June 30 to July 4, 1890, inclusive, for the purpose of instructing the medical department thereof in its duties in camp, and on the completion of this duty will return to his proper station. —Par. 5, S. O. 143, Adjutant-General's Office, Washington, D. C., June 19, 1890.

TAYLOR, MARCUS E., *Captain and Assistant Surgeon*.—Is hereby granted leave of absence for one month, on surgeon's certificate of disability, with permission to go beyond the limits of this Division, and to apply for an extension of five months.—Par. 1, S. O. 45, Division of the Pacific, San Francisco, Cal., June 13, 1890.

INDEX.

- ABBOT, climato-therapy, 67
 Abdominal section, flushing of peritoneum after, 148
 surgery, drainage in, 271
 retrospect of, Price, 646
 tumor, removal of, 189
 viscera, transposition of, 541
 Abducens paralysis, 236
 Abscess complicating uterine fibroid, Davis, 559
 of cornea, Hansell, 252
 of liver, diagnosis of adhesions in, 197, 208
 Accommodation, theory of, 354
 Acid, trichloracetic, in nasal diseases, 541
 Acne, boric acid and alcohol in treatment of, 66
 prescription for, 346
 treatment of, 286
 Acute pancreatitis, 127
 Adhesions in hepatic abscess, diagnosis of, 197
 Adirondack sanitarium, statistics of, 405
 Adulterations with terra alba, 187
 Advertising of proprietary medicines, 488
 Albuminuria in the apparently healthy, Washburn, 356
 significance of, 481
 Alcohol and tobacco headache, prescription for, 231
 in patent medicines, 267
 use of, during lactation, 486
 Alopecia areata, Duhring, 353
 Alvarenga prize, 188
 American Medical Association, proceedings of, 570
 Surgical Association, proceedings of, 547
 Ammonium bromide fumes in asthma, 655
 picrate in malaria, 258
 Amputation in a diabetic subject, Sharpless, 393
 of "tapiroid" cervix, Goodell, 493
 Amylene hydrate in epilepsy, 70
 Amyloid tissues, method of staining, 128
 Anæmia, arsenic and mercury in, 347
 Anæsthesia with bromide of ethyl, 512
 with cocaine, 398
 in litholapaxy, 287
 Anæsthetics, 209
 Anal eczema, treatment of, 101
 fissure, 222
 fistula, case of, 427
 in phthisis, 563
 non-operative treatment of, 258
 Analgesic properties of exalgine, 314
 Anastomosis, intestinal, 185
 Anencephalic monster, 634
 Anatomical material, scarcity of, Roberts, 68
 Aneurism of aorta, MacDonnell, 161
 Angina pectoris, false, 181
 Anomalous diverticulum from ileum, 146
 Anthelmintic, rhubarb as an, 257
 Antidote to snake poison, 59, 55
 Antifebrin, the dose of, 287
 Antimony in inflammation, 398
 Antipyrine and spirit of nitrous ether, 405
 in sea-sickness, 80
 in whooping-cough, 126
 Antiseptic gauze, Lister's new, 147
 irrigation of knee, Weir, 125
 midwifery, Lusk, 581
 paste, 654
 properties of calcium iodate, 370
 surgery, 154
 treatment of erysipelas, 128
 of gastro-enteritis, 287
 of urethritis, White, 637
 of vaccination, 233, 541
 Anus, imperforate, case of, 626
 Aorta, aneurism of, 161
 Aphonia, hysterical, treatment of, 201
 Appendicitis, 549, 605
 quiescent, laparotomy for, 226
 Appendix vermiformis, perforation of, 174
 removal of, Dennis, 698
 Arrhythmia, cardiac, 387
 Aristol, a substitute for iodoform, 205
 Army, examinations for vacancies in, 188
 Arsenic as a cause of multiple neuritis, 102
 in treatment of anæmia, 347
 Arthritis, tubercular, iodoform in treatment of, 126
 Arthrotomy of elbow-joint, 69
 Artificial infant-feeding, a reform in, 290
 Aseptic catgut, 398
 sponges, method of preparing, 19
 urethral instruments, 656
 Ashurst, hospital notes, 255
 Asphyxiation in infants, method of treating, 186, 287
 Aspiration of knee-joint, 309
 Association of American Physicians, proceedings of, 547, 564
 of Medical Editors, 579
 Asthma as a neurosis, Carpenter, 13
 oil-silk jacket in treatment of, 415
 thyroid, 594
 treatment of, 49, 233, 371
 with ammonium bromide fumes, 655
 Astragalus, removal of for equino-varus, 23
 Asylums for insane in New York, 108
 Atropine instilled into eye, hallucinations from, 369
 poisoning, 79
 Aulde, antipyrine and spirit of nitrous ether, 405
 Auscultatory percussion, 49
 Axis-traction forceps, Hoffman, 659
 BACILLUS coli communis, relations to Eberth's bacillus, 315
 Bacteriological examinations at autopsies, 185
 large, 596
 Baldness, treatment of, 49
 Baldy, ectopic gestation, case of, 169
 electricity in pelvic disease, 302, 661
 Balanitis from calomel, 288
 Bandages of magnesia-isinglass, 511
 Barr, intermittent fever, case of, 677
 Beaver, peritoneal adhesions in hepatic abscess, 197
 Beck, legal aspects of vivisection, 280
 Belladonna, glycerite of, 205
 Belt, hallucinations from atropine, 369
 Benzoate of bismuth in treatment of chancre, 180
 Bichloride of mercury in anæmia, 347
 in dysentery, 347
 Bigelow, gynecological electro-therapeutics, 499
 Biliary colic, salicylate of sodium in, 456
 Billings, J. S., vital and medical statistics, 1
 Biniodide of mercury in scarlatina, 232
 Bi-polar faradization in gynecology, Smith, 93
 Bismuth benzoate in treatment of chancroids, 180
 subnitrate, incompatibilities of, 182
 Bladder, fibroid tumors of, 593
 irritable, prescription for, 682
 villous tumors of, 244
 Blepharitis, management of, 143
 Blindness, hysterical, in males, 33
 Blood corpuscles, effects of heat on, 399, 435
 pressures, extreme, 58
 syphilitic, effects of mercury on, 535
 Boiled milk, nutritive value of, 100
 Bone, implantation of, 125
 Boric acid and alcohol in treatment of acne, Post, 66
 Bougies weighted with mercury, 655
 Bozeman, irrigating catheter, 320
 Brain surgery, a failure in, 134
 Bright's disease, general treatment of, 456
 Bridge, appendicitis, 549
 Broad ligament, varicose veins in, 538
 Bromide of ethyl anæsthesia, 512
 of rubidium in epilepsy, 72
 Bronchitis, eucalyptus in, 561
 chronic treatment of, 654
 prescription for, 682
 Brower, types of insanity, 412
 Bryonia alba as a hæmostatic, 512
 Brush, Cheyne-Stokes respiration, 592
 Buboes, treatment of, 47, 233
 Buckmaster, treatment of fibro-myomata by electricity, 91
 Bulkley, dangers of syphilis in dentistry, 449
 Bullous dermatosis, case of, 630
 Burns, treatment of, 71, 182, 370
 Bussey, oil-silk jacket in the treatment of asthma, 415
 CÆSAREAN section, case of, 333
 post-mortem, substitute for, 561
 technique of, 480
 Caffee, 476
 Calcium iodate as an antiseptic, 370
 Calculus, vesical, in boys, 530
 large, 596
 Calf pepsin, 17
 Calomel as a cause of balanitis, 288
 in phthisis, 182
 vaporized, inhalations of, in croup, 455
 Camphoric acid in night-sweats, 481
 Cannabis indica poisoning, Glover, 478
 therapeutics of, 429
 Cancer of intestine, triple resection for, 682
 mammary, extirpation of, 628
 pyloric, gastro-enterostomy for, 112
 rectal, colotomy for, 31
 secondary, suppuration in, 515

- Cancer, uterine, method of treating, 63
 Capsicum as a counter-irritant, 684
 in pneumonia, 313
 Capsules, quinine, short-weight, 487
 Carbolic acid gangrene, 385, 519
 injections in tetanus, 542
 Carbonic acid in treatment of phthisis, 372
 Carbuncles and furuncles, treatment of with iodine, 209
 Cardiac arrhythmia, etc., treatment of, Patton, 387
 affections of menopause, 371
 Caries of teeth, prevention of, 206
 Carpenter, asthma as a neurosis, 13
 Carilage, loose in knee-joint, removal of, 427
 Cartwright lectures, 1
 Casselberry, hysterical aphonia, treatment of, 201
 Castration, double, 479
 Catappa speciosa, Schneek, 624
 Cataphoresis in treatment of syphilitic growths, 675
 Cataract operations, Pipino, 554
 Catarrh of stomach, treatment of, 231, 233
 Catarrhal fever, 133
 Catgut, method of rendering aseptic, 398
 Catheter, self-exhausting irrigating, 320
 Catheterization, dangers of, 456
 Cauda equina, lesions of, 217
 Caustic paste, formula for, 208
 Cautery in treatment of epilepsy, 232
 Cerebro-spinal meningitis, iodoform in, 372
 Cerna, cases treated with iodol, 241
 Cervical paraplegia from dislocation, 236
 Cervix, "tapiroid," amputation of, 493
 detachment of during labor, Hirst, 653
 Chancres, difficulties in diagnosis of, 631
 of breast, 632
 Chancroids, bismuth benzoate in, 180
 Chapped hands, prescription for, 370, 682
 Chilblains, prevention of, 231
 treatment of, 429
 Chloral, administration of, in convulsions, 49
 suppositories of, 50
 Chloralamid, effects of, 47, 595
 Chloroform, after-effects of, 429
 cause of death from, 190
 Chlorosis, treatment of, 103, 314, 682
 Choice of methods in treatment of uterine cancer, Jackson, 63
 Cheyne-Stokes respiration, 589, 592
 Chorea, etc., ocular treatment of, 52
 sulphonal in, 275
 Chronic osteomyelitis, operations for, 211
 prostatitis, treatment of, 150
 rheumatic sore throat, Ingals, 266
 Cinchonidine salicylate, 442
 Climatic treatment of gastro-intestinal disease in children, 39
 of phthisis, 472
 Climo-therapy, individuality the basis of, Abbot, 67
 Clinical aspects of influenza, Davis and Van-neman, 45
 Cocaine anaesthesia in litholapaxy, 287
 in surgery, 398
 Cocillana, uses of, 71
 Codeine in ovarian pain, 149
 therapeutics of, 482, 710
 Cod-liver oil emulsion, Graham, 283
 Co-education, 73, 75, 77
 Coffee, oil of, 476
 Cohen, eucalyptol in bronchitis, 561
 ichthyol ointment in pharyngitis, 178
 terpin hydrate and salol in "colds," 147
 Collodion in atony of tympanic membrane, 314
 in treatment of naevus, 128
 salolized, for rheumatism, 343
 Colotomy for rectal cancer, 31
 stricture, 437
 Compound fractures, treatment of, Dennis, 416
 Congenital abducens paralysis, etc., 236
 stricture of rectum, Kelsey, 224
 Constipation, habitual, treatment of, 543
 in women, treatment of, 542
 Contagiousness of influenza, Trudeau, 185
 Continued fevers of the South, Johnston, 651
 Contusion of intestines, laparotomy for, 227
 Convulsions, infantile, enema for, 50,
 treatment with chloral, 49
 Cornea, abscess of, 252
 Corneal lesions, fluorescein in diagnosis of, 482
 ulcers, eserine in, 336
 Corpus luteum in non-pregnant women, Hirst, 358
 Corpuscles, effects of heat on, 399, 435
 Correspondence—
 Ann Arbor, 25
 Baltimore, 239, 518
 Boston, 404
 Cincinnati, 214, 691
 Detroit, 24, 185
 Havana, 517
 London, 105, 295, 434, 658, 688
 Louisville, 376
 Milwaukee, 238, 379
 Munich, 107, 376
 New Orleans, 462, 635, 690
 New York, 74, 213, 322, 433, 491, 659
 Paris, 237, 292, 377, 463
 St. Louis, 212, 321, 350, 516
 Toledo, 266
 Vienna, 215, 294, 321, 489, 632, 690
 Corrosive sublimate in granular lids, 232
 in dysentery, 347
 Coryza, acute, treatment of, 288
 insufflation for, 50
 menthol in treatment of, 207
 Craniotomy, case of, 340
 Creasote, formula for, 542
 in chronic gonorrhoea, 180
 in phthisis, 50
 intra-tracheal injections of, in phthisis, 289
 parenchymatous injection of, in phthisis, 287
 Crematory, a new, 79
 Creolin, impurities of, 268
 in gynecology, 47
 pills, 128
 Croup and diphtheria, treatment of, 50
 inhalations of vaporized calomel in, 455
 Cumberland, Md., epidemic of typhoid, 401, 403
 Cyanide dressing, Lister's, 130
 gauze, preparation of, 147
 Cystitis, treatment of, 102
 Cystotomy, suprapubic, 224, 322
 DA COSTA, influenza, 60
 Daly, wakefulness of neurasthenia, 360
 Davis, digestion in infants, 146
 influenza, 45
 puerperal fever, 199
 uterine fibroid, case of, 559
 Deafness, pilocarpine in, 655
 Death from chloroform, Wood and Hare, 190
 Deaver, Leiter's urethroscope, 667
 oesophagotomy, 475
 trephining for extra-dural hæmorrhage, 176
 Degeneration of sebaceous glands, Dühring, 353
 Delirium tremens, treatment of, 346
 Delivery by uterine expression, 183
 Dengue fever, 20
 Dennis, removal of appendix, etc., 698
 treatment of compound fractures, 416
 Dentistry, dangers of syphilis in, 449
 Dentition, management of, 542
 Depilatory paste, 542
 Dermoid cyst, complicated by a twisted pedicle, Nelson, 465
 De Saussure, filaria, 704
 De Schweinitz, eserine in corneal ulcers, 336
 ocular complications of malaria, 609
 Diabetes, amputation during, 393
 Diagnosis of adhesions in hepatic abscess, 197
 of syphilis and malignant disease, 148
 use of the hypodermic syringe in, 149
 Diagnostic signs of melancholia, 149
 Diarrhoea, infantile, treatment of, 48
 Diathesis, uric acid, treatment of, 180
 Diazon reaction, 128
 Diet in disease of the kidney, 285
 Digestion in infants, preliminary note on, Davis, 146
 Digitalis and veratrum viride in treatment of pneumonia, 57
 therapeutics of, 109
 Diller, diverticulum from ileum, 146
 Diphtheria, antiseptic application for, 512
 and croup, treatment of, 50
 sulphur in treatment of, 150, 255
 and quinine in treatment of, 709
 Diplomas, American, value of, in Berlin, 240
 Diseases of uterine appendages, treatment of, 457
 Disinfectant injection, 71
 Dislocation, cervical, causing paraplegia, 236
 of the humerus, subcoracoid, etc., Graham, 308
 of lens, 354
 Dissecting material, scarcity of, 68
 Diverticulum from ileum, Diller, 146
 Dobell's treatment of influenza, 70
 Dodge, imperforate anus, 626
 Doses of digitalis and strychnine, 149
 Downs, Cheyne Stokes respiration, 589
 Drainage, defective, as a cause of pharyngitis, 128
 in abdominal and pelvic surgery, Price, 271
 Drake, sulphur in diphtheria, 255
 Dropsy, management of, Tyson, 671
 treatment of, 562
 Dühring, clinical lecture, 353
 Duties of mothers in nursing their children, 155
 Dysentery, complicating intermittent fever, 677
 corrosive sublimate in, 347
 salicylic acid in, 562
 treatment of, 655
 Dysmenorrhoea, excessive, 383
 membranous, 471
 prescription for, 257
 treatment of, 344
 Dyspepsia, formula for, 234
 EAR, labyrinth of, 494
 prominent, operation for, 181
 Eberth's bacillus, relations of, to bacillus coli communis, 315
 Ecthyma, infantile, 632
 Ectopic gestation without characteristic symptoms, Baldy, 169
 Eczema, Dühring, 353
 of anus, treatment of, 101, 288
 rubrum, 283
 Taylor, 229, 680
 treatment of, 206, 288, 346, 709
 Education of women, higher, 75, 77
 Ehrlich's reaction, 128
 Elbow-joint, arthrotomy of, 69
 Electrical execution, White, 458
 treatment of warts, 346
 Electricity, execution by, 79
 in fecal impaction, etc., Hinsdale, 250
 in gynecology, 82, 84, 88, 91, 93, 95, 98, 99, 104
 in pelvic inflammatory diseases, Baldy, 302
 in treatment of fibro-myomata, Buckmaster, 91
 of syphilitic growths, Woodbury, 675

- Electricity in treatment of uterine fibromata, Skene, 99
- Electrolysis in treatment of stricures, 74
- Electro-therapeutics, gynecological, 499
- Emulsion of cod-liver oil, 283
- Empyema, 483
- Jacobi, 117, 136, 170
- Empyreumatic oil of coffee, Reichert, 476
- Endometritis, treatment of, 594, 628
- Enema for infantile convulsions, 50
- nutritive, 313
- Enteric fever, 616
- Enucleation of uterine fibroid, Goodell, 493
- Epidemic influenza, Tyson, 133
- of typhoid at Cumberland, Md., 401, 403
- Epilepsy, amylene hydrate in treatment of, 70
- focal removal of hand-centre in case of, 381
- ocular treatment of, 52
- prescription for, 346
- rubidium bromide in, 72
- treatment of with actual cautery, 232
- trephining for, 253, 479
- Epithelioma, caustic paste for removal of, 208
- treatment of, 632
- Equino-varus, removal of astragalus for, 23
- Ergotin, preparation of, for hypodermic use, 208
- Erysipelas, application for, 257
- Erythema, antiseptic treatment of, 128
- significance of, 71
- Eserine in corneal abscesses, Hansell, 252
- ulcers, de Schweinitz, 336
- Essential paroxysmal tachycardia, Robison, 340
- Ether enemata in treatment of intestinal obstruction, 50
- Ethyl bromide anaesthesia, 512
- Eucalyptus in bronchitis of children, Cohen, 561
- Evolution of venereal diseases, 156
- Exalgine, analgesic properties of, 314
- tests for, 343
- Examinations for appointment in the U. S. army, 188
- Excision of hip-joint, 69
- of mastoid cells, 541
- of varicose veins, 179
- Execution by electricity, 79, 458, 519
- Exophthalmic goitre, treatment of, 429
- Exostosis of femur, removal of, 479
- Expectorant, gurjun oil as an, 628
- Expert testimony, etc., 337
- External urethrotomy, 69
- Extirpation of benign thyroid tumors, 150
- Extra-uterine pregnancy, case of, 169
- Extreme blood pressures, Mitchell, 58
- Eyes, massage of, 101
- Eye-strain, method for detecting, 349
- FACE presentation, complicated by spinal tumor, 340
- Facial paralysis, etc., 236
- Faecal impaction, electricity in, 250
- Faeces, solvents for, 250
- Faradization in gynecology, 93
- Feet, sweating of, 19, 50
- Femur, intra-capsular fracture of, sutured, 126
- Fever, catarrhal, 133
- hepatic, 325
- puerperal, 199
- urethral, prevention of, 345
- Fevers, continued, of the South, 651
- management of, 154
- Fibroid tumor complicating labor, Caesarean section, 333
- of uterus, complicated, 559
- enucleation of, 493
- Fibromata, electricity in treatment of, 99
- uterine, galvanic treatment of, 88
- Fibro-myomata, electricity in treatment of, 91
- Filaria sanguinis hominis, 656
- De Saussure, 704
- Filters, inefficiency of, 511
- Fisher, pneumonia, etc., 539
- Fissure of anus, Kelsey, 222
- Fistula, anal, non-operative treatment of, 258
- case of, 427
- in phthisis, 563
- Floating cartilage in knee-joint, removal of, 427
- Fluorescein in ophthalmic diagnosis, 482
- Flushing the peritoneum, 148
- Focal epilepsy, trephining for, 381
- Foetuin, labor complicated by measles, etc., 254
- Foetus, influence of maternal haemorrhage on, 657
- Forceps, axis-traction, plea for, Hoffman, 649
- Foreign bodies in larynx, treatment of, 286
- Fracture of patella, results of treatment, 318
- of radius and ulna, 23
- compound comminuted, of forearm, 308
- Fractures, compound, treatment of, 416
- France, medical profession in, 188
- French Society of Dermatology and Syphilography, 630
- Furuncles and carbuncles, treatment of, with iodine, 209
- Fussell, tetanus in infants, 145
- GALL-STONE colic, salicylate of sodium in, 456
- Galvanic battery, a new portable, Martin, 95
- treatment of uterine fibromata, McGinnis, 88
- Galvano-chemical cauterization in metritis, 98
- puncture in treatment of pelvic tumors, 84
- Gangrene, carbolic acid, 385, 519
- of the intestine, treatment of, Penrose, 277
- Gastric catarrh, acute, treatment of, 231
- ulcer, A. H. Smith, 526
- Gastro-enteritis, antiseptic treatment of, 287
- enterostomy and pylorectomy, Tuholske, 503
- for cancerous obstruction of pylorus, Stamm, 112
- intestinal disease, climatic treatment of, 39
- Gauze, antiseptic, Lister's new, 147
- Gelatins, Unna's, medicated, 396
- General paralysis, early symptoms of, 455
- Genu-pectoral position in gynecology, Martin, 204
- Germany, medical examinations in, 240
- Gibb, Caesarean section, 333
- Gibney, tubercular ulceration of intestine in hip disease, 306
- Gleet, treatment of, 430
- Globus hystericus, caused by laryngeal tuberculosis, 430
- Glover, cannabis indica poisoning, 478
- Glycerite of belladonna, 205
- of calf-pepsin, Woodbury, 17
- Goitre, enucleation of, Keene, 693
- exophthalmic, treatment of, 429
- Goelet, galvano-puncture in treatment of pelvic tumors, 84
- Gonococcus, non-specific character of, 632
- Gonorrhoea, abortive treatment of, 684
- antiseptic treatment of, 637
- chronic, creasote in, 180
- in a brother and sister, 541
- prescription for, 19, 258, 371
- treatment of with ointments, 155
- with salol, 49, 456
- Goodell, clinical lecture, 493
- dysmenorrhoea; ovaritis, 383
- ovariotomy, 521
- removal of abdominal tumor, 189
- Gout, prescriptions for, 542
- Gradle, blepharitis, 143
- Graduated tenotomy, report on, 51, 52
- Graham, cod-liver oil emulsion, 283
- dislocation of humerus and fracture of forearm, 308
- Granular lids, corrosive sublimate in, 242
- Graves's disease, treatment of, 429
- Gurjun oil as an expectorant, 628
- Gynecological electro-therapeutics, Bigelow, 499
- treatment, abuse of, 457
- work and methods, 5
- HABITUAL constipation, treatment of, 543
- Haematoma of cervix uteri, Kempf, 369
- Haematozoon, malarial, 183
- Haemorrhage after tonsillotomy, 231
- extra-dural, trephining for, 176
- post-partum, treatment of, 206, 344
- symptomatic treatment of, 151
- uterine, hydrastinin in, 395
- in pregnancy, 595
- Haemorrhoids, treatment of, 562
- Haemostatic properties of bryonia, 512
- Hallucinations from atropine in eye, 369
- Hansell, abscess of cornea, 252
- Harlan, dislocation of lens, etc., 354
- hysterical blindness in males, 33
- Hare and Wood, cause of death from chloroform, 190
- Headache from alcohol and tobacco, prescription for, 231
- Heart, arrhythmia, etc., 387
- Hemenway, medical portion of pension bureau, 122
- Hemiplegia, treatment of, 397
- Hepatic abscess, diagnosis of adhesions in, 197
- treatment of, 208
- fever, Pepper, 325
- Hernia, inguinal, in child, operation for, 427
- operation for radical cure of, 81, 179, 286
- Herpes, abortive treatment of, 512
- menstrualis, 343
- progenitalis, Lydston, 141
- zoster of mouth, 676
- Hiller, effects of mercury, etc., 535
- Hinsdale, electricity in faecal impaction, 250
- Hip-disease and general tuberculosis, 306
- treatment of, 361
- Hip-joint, excision of, 69
- Hirst, corpus luteum in non-pregnant women, 358
- detachment of cervix in labor, 660
- substitute for post-mortem Caesarean section, 561
- varicose veins in broad ligament, 538
- History of ovariectomy in America, McMurtry, 272
- Hoagland laboratory, 28
- Hodgen suspension-splint, Mudd, 506
- Hoegh, trephining for epilepsy, 253
- Hoffman, axis-traction forceps, 649
- Homeopathic pharmacy, curiosities of, 660
- Homeopathy, defence of, 264
- in New York, 27
- Hospital notes, Ashhurst, 225
- McBurney, 342, 427, 479
- Nancrede, 69, 309
- Parkes, 311
- Taylor, 229, 283, 393
- Weir, 125, 179
- Hugenschmidt, herpes zoster of mouth, 676
- Humerus, dislocation of, subcoracoid, etc., 308
- Hutchinson, Dr. James H., resolutions upon death of, 108
- W., naso-pharynx as an index of disease, 556
- Hydrarthrosis, puncture and injection in, 258
- Hydrastinin in uterine haemorrhage, 395
- Hydrastis canadensis, 542
- Hydranaphthol in treatment of tinea tonsurans, 18

- Hydrophobia from bite of a cat, 160
 Hygienic management of the newborn, 156
 Hypnotic, prescription for a, 49
 Hypodermic administration of iron, 20
 injections in syphilis, 285
 syringe in diagnosis, 149
 use of ergotin, 208
 Hystero-epilepsy, oöphorectomy for, 465
 Hysterical aphonia, etc., treatment by cauterization of turbinated, 201
 Hysterectomy, case of, 309
 intestinal obstruction after, 597
 supra vaginal, Kelly, 695
 Hysterical blindness in males, Harlan, 33
- ICHTHYOL ointment in pharyngitis, Cohen, 178
 uses of, 428
 Ileum, diverticulum from, 147
 Immersion-treatment of typhoid fever, 480
 Imperforate anus, Dodge, 626
 Implantation of bone, 125
 of mucous membrane for traumatic urethral stricture, 125
 Incontinence of urine, treatment of, 398, 596
 Incompatibilities of bismuth subnitrate, 182
 India, deaths from snakes in, in 1888, 240
 Indian hemp, uses of, 429
 medicine, 323
 Induction of premature labor, 256
 Indurating oedema, 393
 Infant feeding, reform in, 290
 sudden death of, from unusual cause, 292
 Infantile diarrhoea, treatment of, 48
 digestion, 146
 tetanus, 145
 Infants, young, hygienic management of, 156
 Inflammation of appendix, Bridge, 549
 Influenza, 133, 261
 clinical aspects of, 45
 contagiousness of, 185
 Dobell's treatment of, 70
 insurance losses, 407
 menthol in treatment of, 207
 notes on, 78
 the epidemic of, Da Costa, 60
 treatment of, 20, 26
 Ingals, rheumatic sore throat, 269
 Inguinal colotomy for cancer of rectum, Wharton, 31
 hernia in child, operation for, 427
 Inhalation of vaporized calomel in croup, 455
 Injection, disinfectant, 71
 for gonorrhoea, 19
 in hydrarthrosis, 258
 intra-tracheal, of creasote in phthisis, 289
 Innervation of larynx, 71
 Insane asylums in New York, 108
 in Canada, treatment of, 132
 Insanity from traumatism, 101
 ovarian, laparotomy for, 521
 types of, Brower, 412
 Insurance losses from influenza, 407
 medical examinations for, 372
 Intermediate laparotomy for appendicitis, Weir, 226
 Intermittent fever, case of, 677
 International Medical Congress, Tenth, rules of, 159
 Intestinal anastomosis, 185
 antiseptics, 629
 diseases of infants, 512
 obstruction after hysterectomy, 597
 enemata of ether in treatment of, 50
 surgery, Shimwell, 389
 Intestine, gangrene of, treatment, 277
 triple resection of, 682
 tubercular ulceration of, 306
 Intra-capsular fracture of femur, suturing of, 126
 -nasal treatment, abuse of, 258
- Intra-tracheal injections of creasote in phthisis, 289
 -venous injections of quinine in malaria, 100, 232
 Intubation versus tracheotomy, 102
 Iodate of calcium as an antiseptic, 370
 Iodides, dangers of in acute nephritis, 430
 Iodinated muslin, 234
 Iodine in treatment of furuncles and carbuncles, 209
 in vomiting, 231
 Iodoform, aristol, a substitute for, 205
 in cerebro-spinal meningitis, 372
 injections in tubercular arthritis, 126
 in surgery, 180
 poisoning, diagnosis of, 631
 -vaseline in treatment of buboes, 47
 Iodol, cases treated with, Cerna, 241
 in syphilis, 511
 therapeutic use of, 260
 Iron, subcutaneous administration of, 20
 Irrigating self-exhausting catheter, Bozeman, 320
- JACKSON, E., scissors for tenotomy, 184
 Jackson, Reeves, uterine cancer, treatment of, 63
 Jackson, W. R., fibroid polyp of bladder, 593
 Jacobi, M. P., empyema, 117, 136, 170
 Jaundice during pregnancy, 102
 Jeffries, sulphonal in chorea, 275
 Johnston, continued fevers of the South, 651
 Jones, villous tumors of the bladder, 244
- KEEN, nephrectomy, 467
 trephining for epilepsy, 381
 Kelly, supra-vaginal hysterectomy, 695
 Kelsey, colotomy for rectal stricture, 437
 stricture of rectum, 222
 Kempf, hæmatoma of cervix, 369
 Kidney, diet in disease of, 285
 King, laparotomy for contusion of intestines, 227
 Knapp, Dr. Herman, suit against, 267
 Knee-chest position in gynecology, 204
 -joint aspiration of, 309
 irrigation of, 125
 floating cartilage, removal from, 427
 Kolipinski, nitro-glycerin poisoning, 540
- LABOR, complicated, Cæsarean section for, Gibb, 333
 premature, induction of, 256
 Labyrinth of ear, Randall, 494
 Lactation, alcoholic and malted drinks in, 486
 influence of menstruation on, 235
 Laparotomy, flushing of peritoneum after, 148
 for appendicitis, 226
 for contusion of intestines, King, 227
 for ovarian cyst, 521
 insanity, 521
 Laryngeal crises in tabes, 236
 tuberculosis, treatment of, 481
 a cause of globus hystericus, 430
 Larynx, motor innervation of, 71
 rheumatism of, 269
 and trachea, foreign bodies in, 286
 Legal aspects of vivisection, Beck, 280
 Legislation relating to medical portion of pension bureau, 122
 Lens, dislocation of, 354
 Leprosy, prevention of, 79
 Lesions of the cauda equina, Mills, 217
 Leucorrhoea in intermittent fever, 677
 Liabilities for injurious patent medicines, 406
 Life insurance, examinations for, 372
 Ligation of subclavian, Wilson, 678
 Lingual hemiatrophy, 631
 Lister's new dressing, 130, 147
 Litholapaxy, cocaine anæsthesia in, 287
 Lithotomy, case of, 679
- Liver, abscess of, peritoneal adhesions in, 197
 treatment of, 208
 Locomotor ataxia, laryngeal crises in, 236
 treatment of, 72
 London, mortality of, in 1889, 267
 Louisville catastrophe, 374
 Lupus erythematosus, 393
 Lusk, antiseptic midwifery, 581
 Lydston, herpes progenitalis, 141
- MACDONNELL, aneurism of aorta, 161
 Magnesia-isinglass bandages, 511
 Malaria, ammonium picrate in, 258
 intra-venous injections of quinine in, treatment of, 100, 232
 mode of administering quinine in, 344
 ocular complication of, 609
 sunflower in treatment of, 233
 Malarial hæmatozoon, 183
 Mammary cancer, extirpation of, 628
 Management of blepharitis, Gradle, 143
 of fevers, 154
 Mariani & Co., letter from, 663
 Martin, E., effects of mercury on blood, etc., 535
 Martin, F. H., electricity in gynecology and a new battery, 95
 Martin, J. N., genu-pectoral position in gynecology, 204
 membranous dysmenorrhœa, 471
 Massage, influence of on secretion of urine, 49
 of the eyes, 101
 Massey, galvano-chemical cauterization, 98
 Mastoid disease, excision in, 541
 Maury, report on gynecological work, 5
 McBurney, hospital notes, 342, 427, 479
 McCall, salicylate of cinchonidine, 442
 McGinnis, galvanic treatment of uterine fibromata, 88
 McGuire, suprapubic cystotomy, 522
 McDermurry, ovariectomy in America, 272
 McDowell Medical Society, proceedings of, 154
 Measles, pneumonia and premature labor, Feetuin, 254
 Medical expert testimony, E. S. Miller, 337
 profession in France, 188
 ignorance, 235
 Society of South Carolina, 149
 Medico-legal Society of New York, 27
 Melancholia, diagnostic signs of, 149
 Membranous dysmenorrhœa, Martin, 471
 Meningitis, iodoform in, 372
 Menopause, cardiac affections of, 371
 Menorrhagia, hydrastinin in, 395
 Menstruation after oöphorectomy, 482
 influence of, on lactation, 235
 profuse, treatment of, 655
 "Mental hemianopsia," 51
 Menthol, intra-laryngeal injections of in phthisis, 47
 in diseases of nose and throat, 207
 in vomiting of pregnancy, 100
 Mercuric chloride in granular lids, 232
 Mercury biniodide in scarlatina, 232
 effects of, on blood of syphilitics, Martin and Hiller, 535
 Merrill, morphine-habit without injury, 307
 Merz Capsule Co., a letter from, 662
 Methacetin, 540
 Metritis, galvano-chemical cauterization in treatment of, 98
 Micro-biology of rheumatism, 657
 Milk, boiled, nutritive value of, 100
 Miller, E. S., expert testimony, 337
 Miller, J. S., face presentation, etc., 340
 Mills, lesions of the cauda equina, 217
 Mitchell, S. Weir, extreme blood pressures, 58
 motor and sensory reinforcement, 297
 Modern treatment of vesical calculus in male children, White, 530
 Molluscum contagiosum, 283

- Morphine-habit without injurious effects, Merrill, 307
treatment of, 344
Morvan's disease, 595
Motor and sensory reinforcement, 297
innervation of larynx, 71
Mucous membrane, implantation of, Weir, 125
Mudd, Hodgen's suspension-splint, 506
Multiple neuritis from arsenical poisoning, 102
Mussanin, an anthelmintic, 314
Myopathy, primary, Sachs, 663
Myopia in schools, 129
Myxœdema, thyroid grafting in, 596
- NEVUS, treatment of, with collodion, 128
Nancrede, hospital notes, 69, 309
Naphthalin, formula for administration of, 20
Naphthol-camphor as a surgical dressing, 47
Naregamia as an expectorant, 594
Nasal treatment, abuse of, 253
Naso-pharynx as an index of disease, Hutchinson, 556
Neilson, stricture of urethra, etc., 707
Nelson, dermoid cyst complicated by a twisted pedicle, 465
Nephrectomy, Keen, 467
Nephritis, acute, dangers of iodides in, 430
Nephro-lithotomy, 206
Neuralgia, insufflation of salt in treatment of, 49
Neurasthenia, wakefulness of, 360
Neurectomy of inferior dental nerve, 179
Neuritis, multiple from arsenic, 102
Neuroses, reflex, 299
New York Academy of Medicine, officers of, 55
charities in 1889, 187
vital statistics for 1889, 492
Night-sweats, treatment of, with camphoric acid, 481
Nitric acid fumes, death from, 408
Nitroglycerin poisoning, Kolipinski, 540
Nose, deformed, reconstruction of, 615
Nurses and their training-schools, 399, 404
Nutritive enema, 313
value of boiled milk, 100
- OBITUARY—
Becker, Otto, 268
Hutchinson, James H., 26
Lawrence, George W., 56
Livingston, William Oswell, 80
MacMillan, Charles, 108
Nichols, Charles H., 26
Prince, David, 27
Sayre, Lewis H., 56
Smith, Henry Hollingsworth, 435
von Troeltsch, Anton, 108
Ocular complications of malaria, de Schweinitz, 609
defects, correction of in treatment of epilepsy, etc., 52
Edema a symptom of pneumonia, 181
indurating, 393
of glottis caused by potassium iodide, 682
Esophagotomy for impaction of artificial teeth, Deaver, 475
Oil of coffee, 476
—silk jacket in treatment of asthma, Busey, 415
Ointments in treatment of gonorrhœa, 155
Oöphorectomy during pregnancy, 465
for ovarian insanity, Goodell, 521
menstruation after, 482
Operation for anomalous abdominal tumor, Goodell, 189
for hernia, 179
for prolapse of rectum, 50
for prominent ears, 181
for radical cure of hernia, new, 286
Operative treatment of hip-disease, Willard, 361
Ophthalmia neonatorum, prevention of, 373
phyctenular or scrofulous, 156
Orexin, 314, 429, 685
Osteitis of hip and general tuberculosis, 306
Osteomyelitis, chronic, operations for, 211
Otitis, suppurative, papain in treatment of, 127
Ovarian pain, codeine in treatment of, 149
tumor, double, 69
Ovariectomy, cases of, 29, 309
history of, 272, 521
Ovaritis, chronic, 383
Oubain in whooping-cough, 562
Oxygen, therapeutics, 533
Ozæna, treatment of, 257, 258
- PANCREAS, extirpation of, 709
Pancreatitis, acute, 127
Papain in suppurative otitis, 127
Paracentesis thoracis, 513
Paraldehyde in treatment of tetanus, 101
Paralysis, general, early symptoms of, 455
removal of spinous processes for, 210
Paraplegia, cervical, 236
Parasiticide ointment, 313
Parker, reconstruction of deformed noses, 615
Parkes, ovariectomy, 29
Patella, fractured, results of treatment in, 318
Patent medicines, liabilities for injuries from, 406
Path of motor and sensory reinforcement, Mitchell, 297
Patton, cardiac arrhythmia, treatment of, 387
Pelvic diseases, use of electricity in, 302
surgery, drainage in, 271
tumors, treatment of by galvano-puncture, etc., Goelet, 84
Penrose, gangrene of intestine, 277
Pension bureau, legislation concerning, Hemenway, 122
Pepper, hepatic fever, 325
Pepsin of calf, 17
Percussion, auscultatory, 49
Perforation of appendix vermiformis, etc., 174, 605
Pericarditis, etc., simulating pneumothorax, 539
Peritoneal adhesions in hepatic abscess, 197
Peritoneum, flushing of, after laparotomy, 148
Peritonitis, suppuration from perforation of the appendix, Winslow, 174
treatment of, 655
tubercular, surgical treatment of, 150
Perityphlitic abscess, 605
Peruvian balsam in local tuberculosis, 511
Petroleum in treatment of scabies, 313
Pharmacopœia, convention for revision of, 152, 483, 545
Pharyngitis, application for, 100
from defective drainage, 128
ichthyol ointment in treatment of, 178
salol in treatment of, 231
Pharynx as an index of disease, 556
Phenacetin, effects of, 49
Phlyctenular ophthalmia, 156
Phthisical patients at climatic resorts, von Ruck, 472
Phthisis, calomel in treatment of, 182
creasote in treatment of, 50
inhalation for, 257
in the light of modern research, 402
intra-tracheal injections of creasote in, 289
local treatment with menthol, 48
pregnancy during, 207
prescription for, 346
special hospital for, 685
treatment of, with carbonic acid, 372
with parenchymatous injections of creasote, 287
- Physiological study of infantile digestion, Davis, 146
Picrate of ammonium in malaria, 258
Pigmentation of pregnancy, prescription for, 398
Pilocarpine in deafness, 655
Pipino, cataract operations, 554
Pityriasis, prescription for, 314
Placenta prævia, treatment of, 682
Pleurisy, chronic, 483
Pneumonia, alcoholic, capsicum in, 313
afebrile, in infants, 486
complicating premature labor, 254
œdema as a symptom of, 181
simulating pneumothorax, Fisher, 539
veratrum viridi and digitalis in treatment of, 57
Poison of snakes and its antidote, 52, 55
post mortem imbibition of, 34
Poisoning, arsenical, as a cause of neuritis, 102
atropine, 79
cannabis indica, 478
iodoform, diagnosis of, 631
nitroglycerin, 540
ptomaine, 711
Politics in medicine, 21
Polyclinic Medical Society, 27
Porro-Cesarean operation, technique of, 480
Post, boric acid in treatment of acne, 66
Post-mortem imbibition of poisons, Reese, 34
—partum hæmorrhage, treatment of, 206, 344
Potassium iodide, as a cause of glottic œdema, 682
permanganate in snake bites, 684
Pregnancy during phthisis, treatment of, 207
syphilis, 680
hæmorrhage during, 595
jaundice during, 102
oöphorectomy during, 465
vomiting of, treatment by menthol, 100
Premature labor complicated by measles, etc., 254
induction of, 256
Prevention and treatment of puerperal fever, Davis, 199
Price, abdominal surgery, drainage in, 271
retrospect of, 646
Prize, Alvarenga, of College of Physicians, 188
Prolapse of rectum, treatment of, 50, 371
Proprietary medicine advertisers, 488
Prostatitis, chronic, treatment of, 150
Prurigo, 283
Pruritus, sodium salicylate in, 224
prescription for, 313
vulvæ, prescription for, 50, 512
Pseudo angina pectoris, 181
Psoriasis, Reynolds, 409
treatment of, 19
Ptomaine poisoning, 711
Puerperal fever, prevention and treatment of, 199
Puncture and injection in hydrarthrosis, 258
Pylorectomy, 628
and gastro-enterostomy, 503
Pyloric obstruction, gastro-enterostomy for, 112
- QUININE and sulphur in diphtheria, 709
capsules, short-weight, 487
intra-venous injections of, 100, 232
in treatment of syphilis, 47
mode of administering, 344
- RADICAL operations for hernia, 81, 286
Radius and ulna, fracture of, 23
Randall, labyrinth of ear, 494
Rectal cancer, colotomy for, 31
prolapse, operation for, 50
treatment of, 381
stricture, operation for, 437
Rectum, congenital stricture of, 222

- Red corpuscles, effects of heat on, 399, 435
 Reed, climatic treatment of gastro-intestinal disease, 39
 Reese, post-mortem imbibition of poisons, 34
 Reeves, enteric fever, 616
 Reflex neuroses, Starr, 299
 Reichert, empyreumatic oil of coffee, 476
 Reinforcement, motor and sensory, 297
 Report on gynecological work, Maury, 5
 Resection of intestine, triple, 682
 of knee, 679
 of sternum, 257
 Resorcin in treatment of whooping-cough, 48, 268
 Resuscitation of asphyxiated infants, 287
 Retention of urine, 431
 dangers of catheterization in, 456
 Retinoscopy, 349
 Reviews—
 Agnew, Principles and Practice of Surgery, 432
 Ashton, Essentials of Obstetrics, 317
 Baas, History of Medicine and the Medical Profession, 348
 Bartley, Text-book of Medical Chemistry, 318
 Billings, National Medical Dictionary, 153
 Braithwait's Retrospect of Medicine, 401
 Broadbent, The Pulse, 630
 Clevenger, Spinal Concussion, 375
 Cragin, Essentials of Gynecology, 688
 Crookshank, Vaccination, its History and Pathology, 401
 Demarquay, An Essay on Pneumatology, 401
 Despris, Treatise on Fractures, 376
 Escovar, Estudios Sobre Termometria Clinica, 130
 Flores, History of Medicine in Mexico, 260
 Fullerton, Hand-book of Obstetric Nursing, 31
 Gant, Students' Surgery, 375
 Gould, New Medical Dictionary, 687
 Hartridge, Refraction of the Eye, 686
 Hayem, Du Saug et de ses Alterations Anatomique, 317
 Keating, Cyclopaedia of Diseases of Children, 22
 How to Examine for Life Insurance, 630
 Macalister, Text-book of Human Anatomy, 456
 Maissch, A Manual of Organic Materia Medica, 290
 Massey, Electricity in the Diseases of Women, 316
 Medical Annual and Practitioners' Index, 515
 Norris, Syllabus of Obstetrical Lectures, 515
 Ostrom, Massage and the Swedish Movements, 375
 Potter, Hand-book of Materia Medica and Therapeutics, 316
 Rauch, Medical Education, etc., in United States and Canada, 658
 Roberts, Cure of Crooked and Deformed Noses, 318
 Shoemaker and Aulde, Materia Medica, Pharmacology and Therapeutics, 485
 Skene, Education and Culture as Correlated to Health and Diseases of Women, 290
 Smith, A. H., Diabetes, 317
 Stielwagon, Diseases of the Skin, 687
 Students' Aid Series, Vols. V. and VI., 261
 Tait, Diseases of Women and Abdominal Surgery, 687
 Transactions of Association of American Physicians, 261
 Treves and Lang, German-English Dictionary, 432
 Reviews—
 Turnbull, Artificial Anæsthesia, 74
 Ultzmann, Neuroses of Genito-urinary Tract, 375
 Winckel, Text-book of Obstetrics, 348
 Worcester, Monthly Nursing, 316
 Year-book of Treatment for 1890, 401
 Reynolds, psoriasis, 409
 Rheumatic sore-throat, 269
 Rheumatism, micro-biology of, 657
 prescription for, 285
 Rhubarb as an anthelmintic, 257
 Rickets, causation and treatment of, 182
 prescription for, 343
 Roberts, J. B., remedy for scarcity of dissecting material, 68
 Robison, paroxysmal tachycardia, 440
 Rockwell, electricity in gynecology, 82
 Rubidium bromide in epilepsy, 72
 Ruck von, climatic treatment of phthisis, 472
 SACHS, primary progressive myopathy, 665
 Salicylate of cinchonidine, McCall, 442
 of sodium in biliary colic, 456
 in typhoid fever, Spears, 478
 Salicylic acid in dysentery, 562
 Salol and terpine hydrate in "colds," Cohen, 147
 in treatment of gonorrhœa, 49, 456
 of pharyngitis, 231
 Salolized collodion for rheumatism, 343
 Salpingitis, diagnosis and treatment of, 544
 Salt, insufflation of, in neuralgia, 49
 Sand filters, inefficiency of, 511
 Sanitary convention, 186
 Sarcoma of jaw, Keen, 467
 Scabies, petroleum in treatment of, 313
 Taylor, 229
 Scarlatina, biniodide of mercury in, 232
 cold baths in, 683
 School near-sight, 129
 Schneck, catalpa speciosa, 624
 Sciatica, treatment of, 287
 Scissors for tenotomy, Jackson, 184
 Scoliosis, caused by torticollis, 204
 Scrofulous ophthalmia, 156
 Sea-sickness, antipyrine in, 80
 Sebaceous glands, degeneration of, 353
 Secondary consequences of shock, Todd, 444
 Sensory and motor reinforcement, 297
 Sharp, Tait's operation, etc., 41
 Sharpless, amputation in a diabetic, 393
 Shimwell, intestinal surgery, 389
 Skene, secondary consequences of, 444
 Skene, electricity in treatment of fibromata, 99
 Skin-grafting, Thiersch's method, 427
 Sloughing of gastrocnemius in diabetes, 393
 Smith, A. H., gastric ulcer, 526
 Smith, A. L., faradization in gynecology, 93
 Smith, J. L., treatment of tuberculosis, 163
 Snake-poison and its antidote, 52, 55, 684
 Sodium salicylate for biliary colic, 456
 in pruritus, 224
 in typhoid fever, 478
 Solvents for faces, 250
 Somnal, effects of, 47
 South Carolina Medical Society, 159
 Sozoiodol in treatment of burns, 71
 Spears, sodium salicylate in typhoid fever, 478
 Spinal tumor complicating labor, J. S. Miller, 340
 Spina bifida, excision of, 479
 Spinous processes, removal of, 210
 Spiritus etherus nitrosi, combination of, with antipyrine, 405
 Subclavian artery, ligation of, 678
 Splint, Hodgen's, 506
 Sponges, method of rendering aseptic, 19
 Staining of amyloid tissues, 128
 Stamm, gastro-enterostomy for pyloric obstruction, 112
 Starr, reflex neuroses, 299
 Statistics, vital and medical, 1
 Sternum, resection of, 257
 "Stevens's operation," report on, 52
 Stomach, catarrh of, treatment of, 231, 233
 Street noises, 544
 Stricture, electrolysis in treatment of, 74
 Strictures of urethra, case of, 427
 of rectum, intestinal obstruction, colotomy, Kelsey, 437
 congenital, 222
 traumatic urethral, treatment of, 125
 with reflex symptoms, 707
 Strophanthus, 627
 in infantile diseases, 232
 Sulphonal in chorea, Jeffries, 275
 disadvantages of, 233
 formula for, 206
 Sulphur in treatment of diphtheria, 150, 255
 Summer complaint, 512
 Sunflower in treatment of malaria, 283
 Suppositories of chloral, 50
 Suppuration in secondary carcinoma, 515
 Suprapubic cystotomy, McGuire, 522
 Vander Veer, 224
 Surgery, antiseptic, 154
 iodoform in, 180
 of brain, case of, with failure, 134
 of intestines, 389
 Surgical dressing, naphthol camphor as a, 47
 treatment of hepatic abscess, 208
 of tubercular peritonitis, 150
 Suspension splint, Hodgen's, 506
 Suturing an intra-capsular fracture of femur, 126
 Sweating of the feet, 19, 50
 Sweet spirit of nitre, combination of, with antipyrine, 405
 Syccosis, treatment of, 345
 Symptomatic hæmorrhage, treatment of, 151
 Synovitis of knee, irrigation in treatment of, 125
 Syphilis, dangers of, in dentistry, 449
 diagnosis of, 148
 during pregnancy, 680
 idol in, 511
 local manifestations of, 396
 modes of infection, 683
 papular and ulcerating, 393
 quinine in treatment of, 47
 Taylor, 229
 thymolate of mercury in, 285
 Syphilitic, blood of, effects of mercury on, 535
 TABES dorsalis, treatment of, 72
 laryngeal crisis in, pathology, 236
 Tachycardia, paroxysmal, 440
 Tait's operation, plea for conservatism in employment of, 41
 Tamponing uterus for post-partum hæmorrhage, 206
 "Tapiroid" cervix, amputation of, 493
 Tariff on books, 432
 Tartar emetic in inflammations, 398
 Taylor, hospital notes, 283, 393
 syphilis, scabies and eczema, lecture on, 229
 Teeth, caries of, prevention, 206
 Tenotomy scissors, 184
 Terra alba, adulterations with, 187
 Terpine hydrate and salol in "colds," 147
 Tetanus in infants, cases of, Fussell, 145
 injections of carbolic acid in, 542
 treatment of with paraldehyde, 101
 Thallin in typhoid fever, 709
 Therapeutic oxygen, Wallian, 533
 Therapeutics of digitalis, Wood, 109
 Thiersch's method of skin-grafting, 427
 Thompson, tumors of frontal lobe, 586
 Throat, inflammation of, ichthyol ointment in, 178
 Thymolate of mercury, hypodermically, in syphilis, 285
 Thyroid-asthma, 504
 grafting in myxedema, 596

- Thyroid tumors, extirpation of, 150
Tincture of capsicum in alcoholic pneumonia, 313
Tinea circinata, Duhring, 353
 hydranaphthol in, treatment of, 18
 tonsurs, 127, 393
Tobacco and alcohol headache, prescription for, 231
Todd, secondary consequences of shock, 444
Tonsillitis, treatment of, 19
Tonsillotomy, hæmorrhage after, 231
Toothache, prescription for relief of, 102
Tooth-wash, antiseptic, 287
Torticollis as a cause of scoliosis, Young, 204
Trachea, foreign bodies in, 286
Tracheotomy *versus* intubation, 102
Training schools for nurses, 399, 404
Transposition of abdominal viscera, 541
Traumatic dislocation of lens, etc., Harlan, 354
Traumatism as a cause of insanity, 101
Trephining, cases of, 155
 for extra-dural hæmorrhage, Deaver, 176
 for epilepsy, Hoegh, 253
 Keen, 381, 479
Trichinæ in swine, 101
Trichloracetic acid in nasal disease, 541
Tri-State Sanitary Convention, 186
Trudeau, contagiousness of influenza, 185
Tubercular abscess, Keen, 467
 arthritis, iodoform in treatment of, 126
 laryngitis as a cause of globus hystericus, 430, 481
 peritonitis, case of, 562
 surgical treatment of, 150
 ulceration of intestine, Gibney, 306
Tuberculosis, general, in hip-disease, 306
 in the light of modern research, 402
 local excision of, 562
 Peruvian balsam in, 511
 pulmonary, prescription for, 346
 treatment of, 163
Tuholske, gastro-enterostomy and pylorotomy, 503
Tumor of abdomen, anomalous, 189
Tumors of bladder, 244
 of brain, 586
 of thyroid, extirpation of, 150
Turpentine in typhoid fever, Wood, 243
Tympa-num, colloid in disease of, 314
Typhoid fever, 616
 at Cumberland, Md., 401, 403
 drinking water in relation to, Vaughan, 641
 etiology of, 315
 immersion in treatment of, 480
 sodium salicylate in treatment of, 478
 thallin in treatment of, 709
 turpentine in treatment of, 243
Tyson, influenza, 133
ULCER, gastric, Smith, 526
Unna's medicated gelatins, 396
 treatment of tinea, 127
Uræmia with Cheyne-Stokes respiration, Downs, 529
Urethral fever, prevention of, 345, 397
 stricture, cases of, 427
 implantation of mucous membrane in, 125
 treatment by electrolysis, 74
 with reflex symptoms, 707
Urethritis, antiseptic treatment of, 637
Urethroscope, incandescent lamp, Deaver, 667
Urethrotomy, external, 69
Uric acid diathesis, treatment of, 180
 eruptions, 631
Urine, incontinence of, in children, 596
 influence of massage on, 49
 retention of, 431
Urobilinuria, significance of, 127
Uterine cancer, treatment of, 63
 expression, delivery by, 183
 hæmorrhage, hydrastinin in, 395
Uterus, tamponing the, for post-partum hæmorrhage, 206
VACCINATION, antiseptic 541
 complicated by gangrene, 631
 of newborn infants, 398
Vander Veer, suprapubic cystotomy, 224
Vanneman, influenza, 45
Varicose veins, excision of, 179
 in broad ligament, Hirst, 538
 treatment, by injection, 511
 of vulva, 563
Vaughan, examination of drinking water, 641
Venereal diseases, evolution of, 156
Veratrum viride and digitalis in treatment of pneumonia, 57
Vermiform appendix, perforation of, etc., 174
Vesical calculus in boys, treatment of, 530
Villous tumors of bladder, Jones, 244
Visual accommodation, 354
Vital and medical statistics, 1
 statistics of New York for 1889, 492
"Vita nuova," analysis of, 28
Vivisection, legal aspects of, 280
 value of, etc, 259
Vomiting, iodine in treatment of, 231
 of pregnancy, menthol in, 100
WAKEFULNESS of neurasthenia, etc., Daly, 360
Wallian, oxygen, 533
Warfield, carbolic acid gangrene, 385
Warts, removal of, by electricity, 346
Washburn, albuminuria, 356
Water, bacteriological, etc., examination of, 379, 641
Weil's disease, 47
Weir, hospital notes, 179
Wharton, colotomy for rectal cancer, 31
White, antiseptic treatment of anterior urethritis, 637
 electrical execution, 458
 vesical calculus in male children, 530
Whooping-cough, antipyrine in, 126
 oubain in, 562
 resorcin in treatment of, 48, 268
Willard, operative treatment of hip-disease, 361
Winslow, suppurative peritonitis, etc., 174
"Wolf in sheep's clothing," 458
Women, education of, higher, 75, 77
Wood and Hare, cause of death from chloroform, 190
Wood, H. C., treatment of pneumonia, 57
 therapeutics of digitalis, 109
 turpentine in typhoid fever, 243
Woodbury, calf-pepsin, 17
 cataphoresis in treatment of syphilitic growths, 675
Wool, test for purity of, 482
Wyman, a failure in brain surgery, 134
YOUNG, torticollis as a cause of scoliosis, 204
ZINC chloride in endometritis, 594

183

Bind in

MEDICAL NEWS.

A WEEKLY MEDICAL JOURNAL.

Vol. LVI. No. 26.
Whole No. 911.

SATURDAY, JUNE 28, 1890.

Per Annum, \$4.00.
Per Copy, 10 Cents.

Contents.

ORIGINAL LECTURES.

Enucleation of Goitre. A Clinical Lecture delivered at the Jefferson Medical College Hospital. By W. W. KERN, M.D., of Philadelphia 693

ORIGINAL ARTICLES.

Supra-vaginal Hysterectomy. Hysteromyomectomy with Suspension of the Stump in the Lower Angle of the Abdominal Incision. By HOWARD A. KELLY, M.D., of Baltimore 695

The Propriety of the Removal of the Appendix Vermiformis during the

Intervals of Recurrent Attacks of Appendicitis. By FREDERIC S. DENNIS, M.D., of New York 698

A Clinical History of Twenty-two Cases of Filaria Sanguinis Hominis, seen in Charleston, S. C., from 1886 to May, 1890. By P. G. DE SAUSSURE, M.D., of Charleston, S. C. 704

A Report of Two Cases of Stricture of the Urethra and Chronic Prostatitis Presenting Interesting Reflex Symptoma. By THOMAS R. NEILSON, M.D., of Philadelphia 707

MEDICAL PROGRESS.

Quinine and Sulphur in the Treatment of Diphtheria—Therapeutics of Green Coffee—Thalin in Typhoid Fever—Prescription for Eczema

in Children—Cancer of the Pancreas; Extirpation; Recovery 709

EDITORIAL ARTICLES.

Codein 710

CORRESPONDENCE.

Five Cases of Ptomaine Poisoning . . . 711

NEWS ITEMS.

Meeting of the Fifth District Branch of the New York State Medical Association—The 17th Annual Session of the Mississippi Valley Medical Association—A Medical Bishop—Dr. Virchow—Professor Nussbaum Pensioned—The University of Halle—Obituary—Large Russian Surgical Fees 713

Official List of Changes in the Stations and Duties of Officers Serving in the Medical Department U. S. Army 713

Entered at the Post Office at Philadelphia as Second-Class Matter. Copyright secured.

SEVENTH EDITION

PLAYFAIR'S MIDWIFERY.

A TREATISE ON THE SCIENCE AND PRACTICE OF MIDWIFERY.

—BY—

W. S. PLAYFAIR, M.D., F.R.C.P.

PROFESSOR OF OBSTETRIC MEDICINE IN KING'S COLLEGE, LONDON, ETC.

New (fifth) American, from the seventh English edition. Edited, with additions, by

ROBERT P. HARRIS, M.D. *In one handsome octavo volume of 664 pages, with 207 engravings and 5 plates. Cloth, \$4; leather, \$5.*

PLAYFAIR'S *Midwifery* has for many years been a favorite authority both among obstetric teachers and general practitioners of the obstetric art. A work of this kind having reached a fifth American from the seventh English edition would seem scarcely to require any extended notice. Of previous American editions the matter has been largely rewritten or remodelled, besides many new short notes have been added. For either the student or the old practitioner this work meets all needs; it is full and yet condensed; it is clear and well arranged. —*Obstetric Gazette*, Nov. 1889.

Truly a wonderful book; an epitome of all obstetrical knowledge, full, clear and concise. In thirteen years it has reached

seven editions. It is perhaps the most popular work of its kind ever presented to the profession. Beginning with the anatomy and physiology of the organs concerned, nothing is left unwritten that the practical accoucheur should know. It seems that every conceivable physiological or pathological condition from the moment of conception to the time of complete involution has had the author's patient attention. The plates and illustrations, carefully studied, will teach the science of midwifery. The reader of this book will have before him the very latest and best of obstetric practice, and also of all the coincident troubles connected therewith. —*Southern Practitioner*, Dec. 1889.

Lea Brothers & Co., Publishers, Philadelphia,

706 & 708 Sansom Street.

The only prominent Emulsion of Cod-Liver Oil introduced directly to the medical profession.
It is advertised exclusively in medical journals.

HYDROLEINE.

(HYDRATED OIL.)

Produces rapid increase in Flesh and Strength.

FORMULA.—Each Dose contains:

Pure Cod Liver Oil.....30 m. (drops)	Soda.....1-3 Grains
Distilled Water.....35 "	Salicylic Acid.....1-4 "
Soluble Pancreatin.....5 Grains.	Hydrochloric Acid.....1-30 "

Recommended and Prescribed by
EMINENT PHYSICIANS Everywhere.
It is pleasant to the Taste and
acceptable to the most delicate Stomach.

IT IS ECONOMICAL IN USE AND CERTAIN IN RESULTS.

HYDROLEINE (Hydrated Oil) is not a simple alkaline emulsion of oleum morrhuae, but a hydro-pancreated preparation, containing acids and a small percentage of soda. Pancreatin is the digestive principle of fatty foods, and in the soluble form here used, readily converts the oleaginous material into assimilable matter, a change so necessary to the reparative process in all wasting diseases.

The following are some of the diseases in which **HYDROLEINE** is indicated:

Phthisis. Tuberculosis, Catarrh, Cough, Scrofula, Chlorosis,

General Debility, etc.

TO BRAIN WORKERS of all classes, **HYDROLEINE** is invaluable, supplying as it does, the true brain-food, and being more easily assimilated by the digestive organs than any other emulsion.

The principles upon which this discovery is based have been described in a treatise on "The Digestion and Assimilation of Fats in the Human Body," and "Consumption and Wasting Diseases," by two distinguished London physicians, which will be sent free on application.

SOLD BY DRUGGISTS GENERALLY.

C. N. CRITTENTON,

SOLE AGENT FOR THE UNITED STATES.

115 FULTON STREET, N. Y.

A Sample of Hydroleine will be sent free upon application, to any physician (enclosing business card) in the U. S.

Lentz's Aseptic Compact Operating Set, No. 10.



We have from time to time made improvements to this set and are now making a perfect aseptic set, which offers especial facilities for aseptic precautions; the blades are soldered into hollow German-silver handles, nickel plated, are light so as not to be unwieldy and admit of a firm grasp when operating. The saw is adjusted to the handle on an entirely new principle, being made to separate easily and to facilitate thorough cleansing.

The handle is entirely of metal and fenestrated to overcome unnecessary weight.

Scissors and Forceps having French locks can be separated, and the slide can be easily removed from Artery and Needle Forceps.

Therefore, no opportunity is offered for the lodgment and development of germs.

The entire set is patterned with especial reference to facility in cleansing.

The instruments can be sterilized by placing them in boiling water, without fear of damaging them. Wood or rubber handles will not admit of this procedure. For price, see case A.

The following instruments are put up in either a fine Mahogany or Morocco case, with nickel trimmings, lined with velvet, and has an extra space for Trepine with handle, and Elevator if desired:

- | | | |
|--|--|--|
| One Amputating Knife (6-in blade). | One Finger Knife. | One Hernia Knife. |
| One Sharp Curved Bistury. | One Sharp Curved Bistury. | Two Scalpels. |
| One Tenotome. | One Pair Scissors, curved or flat. | One Tonsillectomy. |
| One Liston's Bone Forceps, with Spring. | One Liston's Bone Forceps, with Spring. | One Saw, 9-in blade. |
| One Artery and Needle Forceps, improved. | One Artery and Needle Forceps, improved. | One Hamacher's Flat Rubber Tourniquet, with Chain. |
| One Hamacher's Flat Rubber Tourniquet, with Chain. | One Hamacher's Flat Rubber Tourniquet, with Chain. | One Hemostatic Forceps. |
| One Director, with Aneurism Needle. | One Director, with Aneurism Needle. | Two Silver Probes. |
| Silk, Wire, Wax and Needles. | Silk, Wire, Wax and Needles. | |

With the sixteen instruments contained in this case, any ordinary operation may be performed. Size, 22 inches long, 4 inches wide, 2 inches high.

A. German-Silver aseptic Handles on Knives and Saw.	834 00
B. Hard Rubber aseptic Handles on Knives and Saw.	29 00
C. Ebony Handles on Knives and Saw (as shown in illustration).	25 00
Either Set, with Trepine and Elevator in addition.	4 00

DISCOUNT 25 PER CENT. TO PHYSICIANS.

Our Catalogue of 260 pages will be sent on receipt of 20 cents or postage.

CHARLES LENTZ & SONS, Manufacturers of Surgical and Orthopaedic Apparatus.

Established 1866.

18 NORTH ELEVENTH STREET. PHILADELPHIA.

A-PHOSPHORIZED CEREBRO-SPINANT

(FRELIGH'S TONIC.)

FORMULA.

Ten minims of the Tonic contain the equivalents (according to the formulæ of the U. S. P. and Dispensatory)

Tinct. Nux Strychnos,	1 minim.	Tinct. Gentian,	½ minim.
" Ignatia Amara,	1 "	" Columbo,	½ "
" Cinchona,	4 "	" Phosphorus, C. P.	1-300 grain.
" Matricaria,	1 "	Aromatics,	2 minims.

Dose, 5 to 10 drops, in two tablespoonfuls of water.

INDICATIONS:—Paralysis, Neurasthenia, Sick and Nervous Headache, Dyspepsia, Epilepsy, Locomotor-Ataxia, Insomnia, Debility of Old Age, and in the treatment of Mental and Nervous Diseases.

A Baltimore physician, whose diploma dates from 1825, says:—"Your combination I find vastly more effective than any tonic I have ever used. It furnishes a most powerful evidence of the vastly increased power of medicament by combination and judicious pharmaceutical preparation."

While an Ohio physician says:—"I have been in the profession since 1841, and must frankly say have never been much inclined to run after strange gods, but this new manipulation and combination pleases me."

One of the most widely known physicians in the country, residing in Washington, says:—"The elegance of the formula, the small dose required, and its potency go far to recommend the Tonic to the profession in that large class of neuroses so common among brain workers in this country."

A well-known physician of Chicago, in practice since 1859, says:—"It will be a revelation to most physicians. I have found it peculiarly adapted to the mentally overworked Public School teachers, as well as to the worn-out business man."

The above, and many similar letters from the profession, can be examined at our office. Over 13,000 physicians in New England and the Eastern Middle States are prescribing the Tonic regularly.

Price, One Dollar per bottle, containing 100 of the average 5-drop doses.

Physician's single sample delivered, charges prepaid, on application. That every physician may be his own judge of its value irrespective of the opinions of others, we make the following **SPECIAL OFFER**: We will send to any physician, delivered, charges prepaid, a receipt of 25 cents, and his card or letter-head, half a dozen physicians' samples, sufficient to test it on as many cases for a week to ten days each.

The Tonic is kept in stock regularly by all the leading wholesale druggists of the country.

As we furnish no samples through the trade, wholesale or retail, for samples, directions, price lists, etc., address

O. WOODRUFF & CO., Manufacturers of Physicians' Specialties, 88 Maiden Lane, New York City.

SPECIAL ADVERTISING RATES will be made to **PHYSICIANS** desiring to **PURCHASE, SELL OR EXCHANGE PRACTICES.**

Address **ADVERTISING DEPARTMENT of**

THE MEDICAL NEWS,

706 & 708 Sansom St.,

PHILADELPHIA.

SVAPNIA OR **PURIFIED OPIUM.**

FOR PHYSICIANS' USE ONLY.

Contains the Anodyne and Soporific Alkaloids, Codeia, Narceia and Morphia.

Excludes the Poisonous and Convulsive Alkaloids, Thebaine, Narcotine and Papaverine.

SVAPNIA has been in steadily increasing use for over twenty years, and whenever used has given great satisfaction.

To **PHYSICIANS OF REPUTE**, not already acquainted with its merits, samples will be mailed on application.

SVAPNIA is made to conform to a uniform standard of Opium of Ten per cent. Morphia strength.

JOHN FARR, Manufacturing Chemist, New York.

C. N. CRITTENTON, Gen'l Agent, 115 Fulton St., N. Y.

To whom all orders for samples must be addressed.

SVAPNIA IS FOR SALE BY DRUGGISTS GENERALLY.

SALESMEN WANTED.

We have vacancies for two or three first-class Medical Book Salesmen, on our new subscription works.

THE NATIONAL MEDICAL DICTIONARY.

THE AMERICAN SYSTEMS OF GYNECOLOGY AND OBSTETRICS.

TAYLOR'S CLINICAL ATLAS OF VENEREAL AND SKIN DISEASES.

PEPPER'S SYSTEM OF MEDICINE.

Medical book salesmen are *not canvassers*. They sell the physician the "tools of his trade."

Physicians desirous of resigning practice for a year or two and who are prepared to devote their entire time and energy to the work, can earn an excellent income. Application for territory and full particulars supplied on application. Address, stating age and experience, **MANAGER, SUBSCRIPTION DEPARTMENT.**

LEA BROTHERS & CO., 706 Sansom Street, Philadelphia.

"Peculiarly adapted to all the requirements of the sick room."—A. L. LOOMIS, M.D., LL.D.

"Satisfactory to me."—D. HAYES AGNEW, M.D., LL.D.

"Compounded scientifically, odorless, yet very efficient."—J. ADAMS ALLEN, M.D., LL.D.

"Destined to take the place of every other disinfectant."—H. TUHOLSKE, M.D.

PLATT'S CHLORIDES

A liquid disinfectant, powerful, prompt and cheap.

A deodorizer, without odor or objectionable feature.

A germicide, at once active and thoroughly reliable.

An antiseptic of tried and proven efficiency.

Supplied in quart bottles only, by Druggists everywhere. Price, 50 Cents.

To any physician, who for any reason may be still unfamiliar with the practical value of Platt's Chlorides, a sample will be sent, by express, prepaid, on request. Address, giving both Post and Express offices.

HENRY B. PLATT,

36 PLATT STREET, NEW YORK.

Standard Medical Works.

FLINT'S PRACTICE OF MEDICINE.—SIXTH EDITION.

A Treatise on the Principles and Practice of Medicine. Designed for the use of Students and Practitioners of Medicine. By AUSTIN FLINT, M.D., LL.D., Professor of the Principles and Practice of Medicine and of Clinical Medicine in Bellevue Hospital Medical College, N. Y. New (sixth) edition, thoroughly revised and rewritten by the Author, assisted by WILLIAM H. WELCH, M.D., Professor of Pathology, Johns Hopkins University, Baltimore, and AUSTIN FLINT, JR., M.D., LL.D., Professor of Physiology, Bellevue Hospital Medical College, N. Y. In one very handsome octavo volume of 1160 pages, with illus. Cloth, \$5.50; leather, \$6.50.

ERICHSEN'S SCIENCE AND ART OF SURGERY.—8TH EDITION.

The Science and Art of Surgery; Being a Treatise on Surgical Injuries, Diseases and Operations. By JOHN E. ERICHSEN, F.R.S., F.R.C.S., Professor of Surgery in University College, London, etc. From the eighth and enlarged English edition. In two large and beautiful octavo volumes of 2316 pages, illus. with 984 engravings on wood. Cloth, \$9; leather, raised bands, \$11.

BLOXAM'S CHEMISTRY.—FIFTH EDITION.

Chemistry, Inorganic and Organic. By CHARLES L. BLOXAM, Professor of Chemistry in King's College, London. New American from the fifth London edition, thoroughly revised and much improved. In one very handsome octavo volume of 727 pages, with 292 illustrations. Cloth, \$2; leather, \$3.

CHAPMAN'S PHYSIOLOGY.

A Treatise on Human Physiology. By HENRY C. CHAPMAN, M.D., Professor of Institutes of Medicine and Medical Jurisprudence in the Jefferson Medical College of Philadelphia. In one handsome octavo volume of 925 pages, with 605 fine engravings. Cloth, \$5.50; leather, \$6.50.

BRUCE'S MATERIA MEDICA AND THERAPEUTICS.

Materia Medica and Therapeutics. An Introduction to Rational Treatment. By J. MITCHELL BRUCE, M.D., F.R.C.P., Physician and Lecturer on Materia Medica and Therapeutics at Charing-Cross Hospital, London. Fourth edition. In one 12mo. volume, 591 pages. Cloth, \$1.50. *Students' Series of Manuals.*

LEA BROTHERS & CO., PUBLISHERS, 708 & 708 Sansom Street, PHILADELPHIA.

An Open Letter to the Medical Profession

THE INFANT FOOD PROBLEM SOLVED.

NEW YORK, May 1, 1890.

The Annual of the Universal Medical Sciences for 1889 says: "A perfect Infant Food is still a desideratum; such a food will probably be evolved in the mind of some manufacturer who understands the physiology of infantile digestion and the chemistry of milk. A substitute for human milk to approximate the latter closely should be made entirely from cow's milk, without the addition of any ingredient not derived from milk."

"But not alone do we demand that these Milk Foods contain the equivalent of the solids in human milk, and especially of the albuminoids derived from milk, but that the latter be gathered with the utmost care from properly fed animals, transported with the least possible jolting to the factory, maintained during its transit at a low temperature, then transferred to an apparatus for sterilization, and immediately after the latter has been accomplished reduced to the dry state, in order to prevent the formation of those organisms which Loeffler, Pasteur, and Lester have found to develop in fluid milk after boiling under an alkaline reaction. If such a preparation be put into air-tight and sterilized jars, all will have been accomplished that can be done to render the food sterile, and thus fulfil the chief indications in the prevention of the most serious gastro-intestinal derangements."

"Such a food, too, would have the advantage of being easily and rapidly prepared by addition of sterilized water, affording an altogether sterilized food."

To the Medical Profession at large, we submit for examination and trial the perfected Milk Food known as **LACTO-PREPARATA**. We claim that **LACTO-PREPARATA** is an ideal Infant Food, and that it fulfills the above requirements in every particular, except the partial substitution of cocoa-butter for unstable milk-fat. This substitution was made by advice of Prof. Attfield, London, who made extensive tests of its food value and digestibility in the London Hospitals for Infants.

LACTO-PREPARATA is made from cow's milk evaporated in vacuo a few hours after it leaves the udder. In order to have the product correspond in composition with breast-milk, sufficient milk-sugar is added to bring up the carbohydrates and reduce the albuminoids to a proper proportion (17 per cent). The casein is partially predigested (30 per cent), and the remaining portion is rendered like human milk in character and digestibility. The ingredients are perfectly sterilized and placed in hermetically sealed cans; the powdering, bolting, and canning are done in an air-tight room, all air entering and leaving this room is forced by a blower through heavy layers of cotton. **LACTO-PREPARATA** is adapted more especially to infants from birth to six months of age; and by the addition of water alone represents almost perfectly human milk in taste, composition, and digestibility.

Another product of our laboratory which has been before the profession for a number of years is **CARNRICK'S SOLUBLE FOOD**, which, as now prepared and perfected, contains 37½ per cent. of the solid constituents of milk, 37½ per cent. of wheat with the starch converted into dextrine and soluble starch, and 25 per cent. additional milk-sugar. For infants over six months of age it is perfect in every respect; for infants younger than this, **LACTO-PREPARATA** is more suitable, although *Soluble Food* has also been used largely from birth with the most satisfactory results.

Samples will be sent prepaid, also pamphlet giving detailed description.

REED & CARNRICK,
NEW YORK.

FELLOWS' HYPO-PHOS-PHITES

(SYR: HYPOPHOS: COMP: FELLOWS)

Contains **The Essential Elements** to the Animal Organization—Potash and Lime.

The **Oxydizing Agents**—Iron and Manganese;

The **Tonics**—Quinine and Strychnine;

And the **Vitalizing Constituent**—Phosphorus,

Combined in the form of a Syrup, with *slight alkaline reaction*.

It **Differs in Effect from all others**, being pleasant to taste, acceptable to the stomach, and harmless under prolonged use.

It has **Sustained a High Reputation** in America and England for efficiency in the treatment of Pulmonary Tuberculosis, Chronic Bronchitis, and other affections of the respiratory organs, and is employed also in various nervous and debilitating diseases with success.

Its Curative Properties are largely attributable to Stimulant, Tonic, and Nutritive qualities, whereby the various organic functions are recruited.

In Cases where innervating constitutional treatment is applied, and tonic treatment is desirable, this preparation will be found to act with safety and satisfaction.

Its action is Prompt; stimulating the appetite and the digestion, it promotes assimilation, and enters directly into the circulation with the food products.

The Prescribed Dose produces a feeling of buoyancy, removing depression or melancholy, and hence is of great value in the Treatment of MENTAL AND NERVOUS AFFECTIONS.

From its exerting a double tonic effect and influencing a healthy flow of the secretions, its use is indicated in a wide range of diseases.

PREPARED BY JAMES I. FELLOWS, CHEMIST,
48 VESEY STREET, NEW YORK.

Circulars and Samples sent to Physicians on application.

BUFFALO LITHIA SPRINGS (VA.) WATER.

ITS USES AFTER SURGICAL OPERATIONS, ETC.

[Reprint from Editorial in April number, 1890, of the Virginia Medical Monthly.]

Hunter McGuire, M. D., LL. D., etc., formerly Professor of Surgery in the Medical College of Virginia, Ex-President of the Southern Surgical and Gynæcological Association, etc., in a recent conversation with us on the subject of Mineral Springs Waters, made special mention of the use of Buffalo Lithia Water after cases of laparotomy and other surgical operations. Feeling assured that his views on this subject would be of interest to the Profession generally, we requested a statement of his opinion, and obtained his permission to quote him as follows:

"RICHMOND, VA., March 24th, 1890.

"*Dear Dr. Edwards*—I have just received your letter of this date. I use Buffalo Lithia Water very freely in my hospital. After every case of laparotomy, I give this water for its diuretic properties, and because the stomach bears it so well—often retaining it when everything is rejected. Indeed, I use it freely after nearly all my surgical operations. It is especially valuable in supra-public cystotomy. Many years' experience in its use only confirms the good opinion I have so often expressed in regard to it.

"Yours very truly,

"HUNTER MCGUIRE."

Water in cases of one dozen half gallon bottles, \$5.00 per case, F. O. B. here.

THOMAS F. GOODE, Proprietor, Buffalo Lithia Springs, Va.

OLD AND STEADY

pharmaceutical chemists, we want to be in communication with physicians not disposed to be all the time trying experiments.

We want you to know our work. We mean to be quick; not too quick; would rather be right.

Somebody has to try experiments. Where would progress come from?

The point is this: When we adopt a formula the experimenting time is supposed to be over. We have better means of getting experiments tried than putting them out as adopted formulæ.

Fld. Ext. Ergot, for example. Fluid extracts are excellent tests of a maker's quality. Ours are uniform. You never find one wrong. We make them right; and never by guess. There is such a thing as learning from experience, if one works systematically; but how if one's products vary a little from making to making?

We have the endorsement of many years of use for these; we offer nothing hastily.

Let us send you a sample and catalogue.

HANCE BROTHERS & WHITE
PHILADELPHIA.

Also New York, Boston, Chicago,
St. Louis and Pittsburgh.

SPECIAL OPPORTUNITIES

NOW IN THE MEDICAL SERVICES

—OF THE—

ARMY, NAVY AND MARINE HOSPITAL.

THESE government positions are of great value to a young physician. They are for life, if so desired; or one can resign at any time he wishes. The salaries are good, increasing with years, irrespective of promotion, and his quarters are given him. He is allowed to practice outside and in that way can materially increase his income. He is furnished books and apparatus, and encouraged in any original investigation he may undertake. Contrast these facts with the difficulties he has to encounter in private practice. For particulars, address

DRS. DAWBARN & SYMONDS,

345 W. Fifty-sixth Street, New York.

COLLEGE OF PHYSICIANS & SURGEONS OF BALTIMORE, MD.

The PRELIMINARY COURSE will begin September 15, 1890.
The REGULAR SESSION will open October 1, 1890, and close April 1, 1891.

For Catalogue containing full information, write to

THOMAS OPIE, M.D., Dean,

N. W. COR. CALVERT AND SARATOGA STS., BALTIMORE, MD.

MEDICO-CHIRURGICAL COLLEGE OF PHILADELPHIA.

Winter Session will begin Wednesday, October 1st, and continue until April 16th. Preliminary Session begins September 8th. Spring Term, April 20th, 1891.

The Curriculum is graded and a preliminary examination and three Annual Winter Sessions are required. Laboratory instruction in Chemistry, Histology, Pathology, Hygiene, Physiology, with Bedside instruction in Medicine, Surgery and Gynecology, is a part of the regular course.

Fees: Matriculation, \$5.00. First and Second years each, \$75.00. Third year, \$100.00. Fourth year free to those in attendance three sessions, to all others, \$100.00.

For announcement or information, apply to

E. E. MONTGOMERY, M. D., 1818 Arch Street, Philadelphia, Pa.

BELLEVUE HOSPITAL MEDICAL COLLEGE CITY OF NEW YORK.

Sessions of 1890-91.

The REGULAR SESSION begins on Wednesday, September 24, 1890, and ends about the middle of March, 1891. During this session, in addition to the regular didactic lectures, two or three hours are daily allotted to clinical instruction. Attendance upon at least two regular courses of lectures is required for graduation.

The SPRING SESSION consists of recitations, clinical lectures and exercises, and didactic lectures on special subjects. This session begins about the middle of March and continues until the middle of June. During this Session, daily recitations in all the departments are held by a corps of Examiners appointed by the Faculty.

The CARNEGIE LABORATORY is open during the collegiate year, for instruction in microscopical examinations of urine, practical demonstrations in medical and surgical pathology, and lessons in normal histology and in pathology, including bacteriology.

For the annual Circular and Catalogue, giving requirements for graduation and other information, address Prof. AUSTIN FLINT, Secretary, Bellevue Hospital Medical College, foot of East 56th Street, New York City.

UNIVERSITY OF MICHIGAN, DEPARTMENT OF MEDICINE AND SURGERY.

Forty-First Year, commencing Wednesday, Oct. 1, 1890.

The Course of Instruction now comprises three collegiate years of nine months each. Attendance compulsory. Examination for admission required. Women admitted.

Fees:—For the first year—Residents of Michigan, \$35; non-residents, \$60. For each subsequent year—Residents of Michigan, \$25; non-residents, \$35. Graduation Fee—for all alike, \$10.

Circular and Catalogue, with full details, sent on application.

WM. A. CAMPBELL, M. D., Secretary, Ann Arbor, Michigan.

NIAGARA UNIVERSITY, MEDICAL DEPARTMENT, BUFFALO, N. Y.

Next Session begins September 22, 1890, and continues seven months. Course of study three years, graded.

Fees.—Matriculation, \$5.00; Lectures, \$75.00; Practical Anatomy, \$10.00; Perpetual Tickets, \$165.00; Graduation, \$25.00. No other fees.

For announcements, etc., address the Secretary,

ALVIN A. HUBBELL, M. D., 212 Franklin Street, Buffalo, N. Y.

University of Pennsylvania.

MEDICAL DEPARTMENT.

The 125TH ANNUAL WINTER SESSION will begin Wednesday, October 1st, 1890, at 12 M., and continue seven months.

The PRELIMINARY SESSION begins September 22d; the SPRING TERM early in May, 1891.


The Curriculum is graded and three annual winter Sessions are required. Practical instruction, including laboratory work in Chemistry, Histology, Osteology and Pathology, with Bed-side Instruction in Medicine, Surgery and Gynecology, are a part of the regular course and without additional expense.

For Catalogue and announcement containing particulars apply to

DR. JAMES TYSON, Dean,

36th and Woodland Avenue, - Philadelphia.

Free.



Queen's Pocket Pupilometer.

We will send, without charge, to any physician who will send us his name and address, our little pocket pupilometer, as illustrated above, for determining the size of the pupil in Ophthalmic Diagnosis. We will also send our 180-page catalogue of *Ophthalmoscopes, Trial Sets, Trial Frames, Test Cards, Spectacles, Eye Glasses, Lorgnettes, &c.*

We will also send our pad of *Prescription Blanks* by which orders for Spectacles or Eye Glasses may be correctly transmitted. We will also send *Queen's Test Types* for general office use.

We will also send our Phototype Card of the best forms of Spectacles and Eye Glasses. This card is 16 x 12, and is the finest illustration of Spectacles and Eye Glasses ever published.

All of the above will be found most useful to any physician who has Ophthalmic work. Remember we send the entire collection *free*, provided you say where you saw this advertisement.

QUEEN & CO.

924 Chestnut Street, - - Philadelphia.

If you are contemplating the purchase of a set of Trial Lenses, do not omit to ask for a description of *Queen's Complete Set*, containing 110 pairs of Mounted Lenses, Trial Frames, &c., in handsome Morocco and Plush-lined case, for \$75.00.

A NEW TABLET HYPODERMIC SYRINGE.

Has Double Fenestrated Barrel, Finger or Cross Bars.
Unscrews at end for Tablets.



No. 312.

ITS CAPACITY. IT HAS TWO NEEDLES OF REFINED STEEL, CAREFULLY TEMPERED.

Put up in a neat Morocco Case, carefully and thoroughly made, compact and durable, with pockets for recipes and papers, also six vials. Easily carried in vest-pocket. Price, \$3.33, less 25 per cent. to Physicians, or \$2.50 NET. Postage, 4 cents.

SEND FOR CIRCULAR OF OUR OTHER HYPODERMIC SYRINGES.
HYPODERMIC SYRINGES OF ALL KINDS PROMPTLY REPAIRED.

Our Hypodermic Syringes have for years had the highest reputation.

CODMAN & SHURTLEFF, MAKERS OF
SURGICAL INSTRUMENTS,
13 & 15 TREMONT STREET, BOSTON, MASS.

Please mention MEDICAL NEWS.

Approved Medical Text-Books.

**** BARNES' SYSTEM OF OBSTETRICS. **** *A System of Obstetric Medicine and Surgery, Theoretical and Clinical.* For the Student and the Practitioner. By Robert Barnes, M. D., Physician to the General Lying-in Hospital, London, and Fancourt Barnes, M. D., Obstetric Physician to St. Thomas' Hospital, London. The Section on Embryology contributed by Professor Milnes Marshall. Octavo, 872 pages, 231 illustrations. Cloth, \$5.00; leather, \$6.00.

**** DRAPER'S MEDICAL PHYSICS. **** *Medical Physics. A Text-book for Students and Practitioners of Medicine.* By John C. Draper, M. D., LL. D., Professor of Chemistry in the University of the City of New York. Octavo, 734 pages, 376 woodcuts, mostly original. Cloth, \$4.00.

**** FRANKLAND & JAPP'S CHEMISTRY. **** *Inorganic Chemistry.* By E. Frankland, D. C. L., F. R. S., Professor of Chemistry in the Normal School of Science, London, and F. R. Japp, F. I. C., Assistant Professor of Chemistry in the Normal School of Science, London. Octavo, 677 pages, 51 woodcuts and 2 lithographic plates. Cloth, \$3.75; leather, \$4.75.

**** COATS' PATHOLOGY. **** *A Treatise on Pathology.* By Joseph Coats, M. D., F. F. P. S., Pathologist to the Glasgow Western Infirmary. Octavo, 829 pages, 339 beautiful illustrations. Cloth, \$5.50; leather, \$6.50.

**** ASHHURST'S SURGERY. NEW EDITION. **** *The Principles and Practice of Surgery.* By John Ashhurst, Jr., M. D., Professor of Clinical Surgery, University of Pennsylvania. New (fifth) edition, enlarged and revised. In one large and handsome octavo volume of 1144 pages, with 642 illustrations. Cloth, \$6.00; leather, \$7.00. *Just ready.*

**** EMMET'S GYNAECOLOGY. **** *The Principles and Practice of Gynaecology; for the use of Students and Practitioners of Medicine.* By Thomas Addis Emmet, M. D., LL. D., Surgeon to the Woman's Hospital, New York, etc. New (third) edition, thoroughly revised. Octavo, 880 pages, 150 illustrations. Cloth, \$5.00; leather, \$6.00; half Russia, \$6.50.

**** SMITH ON CHILDREN. **** *A Treatise on the Diseases of Infancy and Childhood.* By J. Lewis Smith, M. D., Clinical Professor of Diseases of Children in the Bellevue Hospital Medical College, New York. New (sixth) edition, thoroughly revised and rewritten. Octavo, 867 pages, 40 illustrations. Cloth, \$4.50; leather, \$5.50; half Russia, \$6.00. *Just ready.*

FULL CATALOGUE ON APPLICATION TO

LEA BROTHERS & CO., PUBLISHERS, PHILADELPHIA.

L. C. VANUXEM & Co.

General Agents for Pennsylvania, Ohio, Illinois and Delaware, for the

NEW YORK LIFE INSURANCE COMPANY

WILLIAM H. BEERS, President.

Telling Figures from the Forty-fifth Annual Report

(JANUARY 1st, 1890)

TOTAL INCOME, over Twenty-nine Million Dollars.
BENEFITS TO POLICY HOLDERS, over Twelve Million Dollars.
INTEREST INCOME, nearly Five Per Cent. on Average Net Assets.
NEW INSURANCE WRITTEN, over One Hundred and Fifty-one Million Dollars.
ASSETS, over One Hundred and Five Million Dollars.
SURPLUS, by the New York State Standard, over Fifteen and a Half Million Dollars.
INSURANCE IN FORCE, over Four Hundred and Ninety-five Million Dollars.

PROGRESS IN 1889.

INCREASE IN INTEREST.....	\$ 303,653 06
INCREASE IN BENEFITS TO POLICY HOLDERS.....	1,148,051 61
INCREASE IN SURPLUS FOR DIVIDENDS.....	1,716,849 01
INCREASE IN PREMIUMS.....	3,458,330 35
INCREASE IN TOTAL INCOME.....	3,761,983 41
INCREASE IN ASSETS.....	11,573,414 41
INCREASE IN INSURANCE WRITTEN.....	26,099,357 00
INCREASE IN INSURANCE IN FORCE.....	75,715,465 00

GROWTH OF BUSINESS IN PENNSYLVANIA.

PREMIUMS COLLECTED.

1880.....	\$ 198,000
1881.....	242,500
1882.....	316,800
1883.....	401,500
1884.....	470,200
1885.....	596,900
1886.....	835,100
1887.....	1,047,500
1888.....	1,058,000
*1889.....	1,235,100
*Including Pa. Premiums paid In New York.	
\$6,401,600	

The total Premiums Collected in entire agency in 1889 was \$2,600,000.

We offer two forms of policies combining the Non-forfeiting features originated by the New York Life Ins. Co. in 1860:

One for protection only—where a low rate of premium is charged, the protection large and the returns showing a modest profit in cash. The other form—where the premium is large—will show an investment equal to the face of the policy and fifty per cent. of all premiums paid thereon. For prospectus and full details, address

L. C. VANUXEM & CO.

OFFICES:

PHILADELPHIA, PA., 331 & 333 Walnut Street.

CINCINNATI, O., 99 West Fourth Street.

COLUMBUS, O., 20 East Broad Street.

PITTSBURG, PA., 95 Fifth Avenue.

CLEVELAND, O., 42 & 44 Euclid Avenue.

CHICAGO, ILL., Phenix Building.

WILMINGTON, DEL., 3 W. Seventh Street.

THE AMERICAN SYSTEMS OF GYNECOLOGY AND OBSTETRICS

SYSTEMS OF GYNECOLOGY AND OBSTETRICS

IN TREATISES BY AMERICAN AUTHORS.

GYNECOLOGY edited by **MATTHEW D. MANN, A. M., M. D.**

*Professor of Obstetrics and Gynecology in the
Medical Department of the University of Buffalo.*

OBSTETRICS edited by **BARTON COOKE HIRST, M. D.**

*Associate Professor of Obstetrics in the Univer-
sity of Pennsylvania, Philadelphia.*

In four very handsome octavo volumes, containing 3612 pages, 1092 engravings and 8 colored plates. Per volume: Cloth, \$5.00; leather, \$6.00; half Russia, \$7.00.

Complete Work Just Ready.

FOR SALE BY SUBSCRIPTION ONLY.

ADDRESS THE PUBLISHERS. FULL DESCRIPTIVE CIRCULAR FREE ON APPLICATION.

PROFESSIONAL OPINIONS.

From **WILLIAM GOODELL, M. D.,**

*Clinical Professor of Gynecology in the University of Pennsylvania,
Philadelphia.*

"The American System of Obstetrics" is most admirable, containing the best work of our very best men.

From **PAUL F. MUNDE, M. D.,**

*Professor of Gynecology in the New York Polyclinic; Gynecologist to the
Mount Sinai Hospital.*

It gives me much pleasure to say that after a careful perusal, I find it a work of exceptional excellence which reflects great credit on American medicine, and particularly on American gynecologists. It is a real treat to read a book so carefully and instructively written, so elegantly appointed, and so thoroughly discussing the various subjects which it

contains. No American physician need fear to place it on a par with the best work of foreign authors.

From **FORDYCE BARKER, M. D., LL. D.,**

*Professor of Clinical Midwifery and Diseases of Women in the Bellevue
Hospital Medical College, New York.*

I regard the work as very valuable, by far the most so of any on the subjects of which it treats which have appeared from the American press and I think that I may safely add, from any nationality. I am particularly proud of it as a work by American authors.

From **SAMUEL C. BUSEY, M. D.,**

Fellow American Gynecological Society, etc.

The editor and publishers are to be congratulated on the issue of such a comprehensive and valuable work as Hirst's American System of Obstetrics.

PRESS OPINIONS.

From the **Medical News**, August 25, 1888.

This, the companion volume to the "System of Gynecology by American Authors," equals it in the excellence of the subject matter and the perfection of the publishers' art. As a treatise for the use of the practitioner the volume will be found to represent admirably the obstetric science of the day, as exemplified in American practice.

From the **Boston Medical and Surgical Journal**,
January 3, 1889.

The same qualities which were characteristic of the first volume are conspicuous here: the thoroughly practical way in which the various subjects are treated, a wide knowledge and fair appreciation of what has been done in foreign countries, and withal a distinctive American independence of thought and flavor of originality about the whole.

From the **American Journal of the Medical
Sciences**, March, 1889.

Gynecology for all peoples and times has been enriched by the material gathered together in the two volumes that compose the work. It is a good work from the hands of good and earnest men, and as such it will endure and keep its place.

From the **Obstetric Gazette**, August, 1888.

This is a very valuable contribution to the literature of Obstetrics. The editors, contributors and publishers are entitled to most hearty congratulations for the complete kind of work that has appeared.

From the **Journal of the American Medical
Association**, September 8, 1888.

One is at a loss to know what to say for fear that just and merited praise may be mistaken for flattery. The papers of Drs. Engelmann, Martin, Hirst, Jaggard and Reeve are incomparably beyond anything that can be found in obstetrical works.

From the **University Medical Magazine**,
November, 1888.

In substance it is practical. American obstetricians are the most successful whether performing craniotomy with pocket scissors, miles from an assistant, or performing Cæsarian section in a hospital in the metropolis with the most approved technique. Among the most encouraging signs of the present is the determination on the part of intelligent men to know the science of the world, and this places at our disposal the clinical treasures of the older continents.

LEA BROTHERS & CO., PUBLISHERS, 706 & 708 SANSON STREET, PHILADELPHIA.

NOTES ABOUT SOME SEASONABLE REMEDIES FOR PREVALENT SUMMER DISEASES.

Among the diseases that are especially prevalent during the hot weather are the intestinal disturbances: diarrhoea, dysentery, intestinal colic, gastric irritability, etc.

The remedies commended for these affections are manifold, and must necessarily vary with the individual conditions. Simplicity of medicinal treatment in these cases is, we believe, practiced by the most intelligent physicians. As easily available, eligible and convenient remedies, when other than dietetic or hygienic measures are desirable, we may mention the following, supplied by us, which have proved efficient in the experience of many physicians:

Chloranodyne, a preparation of much value as a sedative, anodyne and antispasmodic in disturbances of the digestive tract incident to summer. The formula is an improvement upon the chlorodyne of J. Collis Browne, M.R.C.S.L., which has long been established in favor abroad.

Among intestinal sedatives tablets of bismuth subcarbonate and subnitrate, antacid tablets, soda-mint tablets, Dover powder tablets, pepsin and bismuth tablets, arsenite of copper tablets, and pepsinum purum tablets or pepsin cordial when diarrhoea is dependent on fermentation of undigested food, offer a choice for selection to meet the varying indications present.

Our pepsin cordial presents the ferment in an especially desirable form for administration, being permanent and palatable, as well as possessing in a high degree the proteolytic properties of the gastric juice.

The antiseptic and sedative treatment of intestinal disorders is a deservedly popular one, and in this class of remedies we would remind physicians of the antiseptic yellow oxide of mercury tablets which have proved of so much service in septic forms of dyspepsia, and prophylactic against diarrhoea and dysentery.

For local use mercuric iodide tablets will be found convenient for making solutions of any desired strength, and for purposes of disinfection of excretions, or surroundings, antiseptic liquid or thiocamph may be employed to advantage.

There is another class of remedies, to which we wish to ask especial attention, of much service in inducing emesis in children or adults in which overloading of the stomach or intestines leads to diarrhoea. A most reliable and certain emetic for this condition is normal liquid ipecac (standard two-per-cent. emetine). This is available also in croup incident to the exposure of children to sudden changes of temperature and the out-door life of summer.

For use as an expectorant for coughs and colds, two recent remedies are being largely employed, viz.: cocillana and Goanese ipecac.

Information as to any of these remedies in the form of descriptive circulars and samples for testing when desired, will be furnished to physicians on request.

PARKE, DAVIS & CO.,

DETROIT AND NEW YORK.

10234

